



Knowledge and use of biocultural diversity by Nahua in the Huasteca region of Hidalgo, Mexico

Tania Vianney Gutiérrez-Santillán¹, Ángel Moreno-Fuentes², Arturo Sánchez-González² and Gerardo Sánchez-Rojas^{2*}

ABSTRACT

In the Huasteca region, high biological diversity and diverse ethnic groups converge. The implementation of metrics for biocultural diversity was developed based on metrics used in analyzes of biological diversity. We compared the results of the diversity known and used by two communities of Nahuatl origin established in two types of vegetation (Tropical Semi Evergreen Forest [TSEF] and Mountain Cloud Forest [MCF]). The fieldwork was carried out from January 2011 to December 2012; the ethnobiological information was collected combining the methods: percentual and snowball, additional applying as an ethnographic tool: multiple free lists and semi-structured interviews to 125 informants. The ethnobiological information was analyzed by adapting indexes used in the evaluation of biological diversity and multivariate methods. The informants identified 409 ethnospecies for both communities, although they only correspond to 383 biological species, the several taxonomic groups. The TSEF presents a greater richness of species-ethnospecies concerning the MCF. Eleven categories of use were identified, with the edible category being the most mentioned. In general, local people have extensive knowledge of the biological diversity present in their territory, and there are distinct differences in knowledge between communities established in different ecosystems. However, there is much similarity in knowledge and use of biodiversity, since both populations belong to the same cultural group. We believe that our results show the relevance of using the metrics used in the evaluation of biocultural diversity.

Keywords: Index of Biodiversity; Multitaxonomic; Ethnobiology

1 Instituto de Ecología Aplicada, Universidad Autónoma de Tamaulipas, Av. División del Golfo #356. Col. Libertad. C.P: 87020, Ciudad Victoria Tamaulipas, México.

2 Centro de Investigaciones Biológicas, Universidad Autónoma del Estado de Hidalgo, Apartado Postal 1-69, Plaza Juárez, C.P. 42001, Pachuca Hidalgo, México.

* Corresponding author. * E-mail address: TVGS (tvgutierrez_santillan@yahoo.com), AMF (amoreno@uaeh.edu.mx), ASG (arturosg@uaeh.edu.mx), GSR (gsanchez@uaeh.edu.mx)

INTRODUCTION

A spatial co-occurrence among biological, ecological, environmental, geographical, linguistic and cultural components can be observed around the world; this co-occurrence show that indigenous groups

mostly populate greatest biodiversity areas, this correlation is conceptualized as biocultural diversity (Maffi 2001; 2005; 2007; Loh and Harmon 2005; Harmon and Loh 2010; Stepp et al. 2004; 2005). This biocultural diversity establishes unique social-ecological relationships (Pretty et al.

2009); since it is recognized that each indigenous group appropriates the biodiversity present in their territories (Boege 2008; Toledo 2010).

Although biocultural diversity is recognized, there are semantic and conceptual gaps in the definition, additionally it has been seen that biological diversity and biocultural diversity are not entirely congruent in terms of the factors that generate them (Cardillo et al. 2015). A concept of biocultural diversity should be based on more than merely the sum of its components. For this reason, we contextualize biocultural diversity as “the variety of organisms that are known, named, classified, organized, used, exploited, domesticated and/or manipulated by different human societies; including the social-ecological systems of which this diversity forms a part at various spatiotemporal scales (Gutiérrez-Santillán 2018).

This conceptualization requires us to define the units that make up biocultural diversity, which are called *ethnospecies*. An *ethnospecies* corresponds to a biological species and its cultural identity. Indispensable attributes of an *ethnospecies* are the combination of the taxa itself (species or genus but not any higher category) with the cultural identity associated with the traditional unique name for that taxa as the cognitive basis for the recognition of the organism by human groups (Hunn 2011).

Assessing the richness and diversity known and exploited by human groups, that is, biocultural diversity, helps us to understand the degree of correlation with biodiversity and thus to evaluate other associated phenomena; for example; understand the spatial congruence between species richness and cultural evolution (Turvey and Pettorelli 2014) as well as the

loss of both diversities (Sutherland 2003). The importance of maintaining the balance between nature-culture and how to sustain it is related to the proper development of future projects (Maffi and Woodley 2008), the use of natural resources as a way of connecting with nature (Grasser et al. 2012), the conservation of biocultural diversity as a goal (Hong 2013) and even as a response to the challenges faced by large cities (Elands et al. 2018).

Biocultural diversity should be used when designing priorities in conservation (Dunn 2008), and social-ecological research should be the link between human and natural aspects (Gavin et al. 2015; Saslis-Lagoudakis and Clarke 2013); integrating the new biocultural perspective into conservation biology (Gavin et al. 2015; Gorenflo et al. 2012; Huntington 2013; Luque and Doode 2010; Saslis-Lagoudakis and Clarke 2013; Wolverton et al. 2014). This is supported by successful conservation programs that integrate traditional knowledge and local practices (Huntington 2013; Maffi and Dilts 2014).

Mexico is one of the countries with high biological diversity (Neyra-González and Durand-Smith 1998), as well as cultural diversity (De Ávila 2008), which makes it one of the most propitious countries for ethnobiological research, from different approaches and themes, for example, the traditional classification (Berlin 1973; Berlin et al. 1973; Hunn 1998; 2008; Alcántara-Salinas et al. 2013), the evaluation of cultural importance (Garibay-Orijel et al. 2007).), the domestication of plants (Bye and Linares 1983), among many other examples. However, few studies comprehensively address the traditional knowledge of different taxonomic groups (Aldasoro-Maya 2012; Argueta-Villamar 2008; Cano 1988; Hunn 1998, 2008).

Because in general ethnobiological works have been developed by defined subdisciplines (ethnobotany, ethnomycology, and ethnozoology).

The Huasteca is a region located in central-eastern Mexico, where the territories of several indigenous peoples converge. It is characterized by high biological diversity (Olivier 2008; Ruvalcaba et al. 2004), associated to many different geographic and environmental condition. Various ethnobiological studies have been carried out in this region, focusing mainly on ethnobotany of the Teneek (Alcorn 1981a; 1981b; 1983; 1984), Nahuas (Andrade-Cetto 2009; Hernández 2003), Pames (Cabajal-Esquivel et al. 2012; Torres et al. 2015), Tepehuas (Álvarez 2002; López-Villafranco and Aguilar-Contreras 2010) and Totonacos (Cano 1988). A number of studies have examined the importance of the social-ecological relationship in terms of magical/religious aspects (Montoya 1968; Gallardo-Arias 2004; Piotrowska 2013a; 2013b), traditional medicine (Andrade-Cetto 2009) and the use of edible fungi (Bautista-Nava 2007; Isidoro-Reséndiz 2011).

The few ethnozoological studies in the region have documented knowledge, perception, classification and, zotherapy. The vertebrate groups studied were fish (Montaño et al. 2010; González et al. 2010), reptiles (Penguilly et al. 2010) and birds (Jaimes et al. 2014). Additionally, research in the Huasteca region has integrated ethnobiological data for multiple taxonomic groups; for example, in Hidalgo (Gutiérrez-Santillán 2013; Hernández and Bautista 2011), in Puebla (López del Toro et al. 2009), in San Luis Potosí (Alonso-Castro et al. 2011) and Veracruz (Cano 1988), which has contributed to recognition of the biocultural diversity of the region.

Considering that the Huasteca region is rich in biodiversity and has the presence of one of the most representative ethnic groups in Mexico the Nahuas, the work is approached as a case study on conceptual and methodological aspects of cultural diversity. We proposed the use of the most current metrics for the analysis of biological biodiversity, to propose it as a new approach to the measurement of biocultural diversity. The study was designed with the following objectives: 1) to identify the ethnospecies of plants, fungi, and animals that the local people know and use, in order to document their degree of knowledge about local biodiversity; 2) to compare ethnospecies richness between the different vegetation types, in order to find out whether traditional knowledge is similar within a given cultural group or whether it may differ between different ecosystems; 3) to analyze the ethnospecies detected in the study by means of diversity indexes as an evaluation metric for use in biocultural diversity studies; 4) to evaluate the degree of correlation between taxonomic species and ethnospecies, in order to find the degree of culturally recognized biodiversity; and 5) to identify use categories, in order to find the association between taxonomic groups and use categories.

We expected to contribute to the subject of biocultural diversity by establishing ethnospecies as social-ecological units of study and their evaluation by using the most recent biodiversity metrics (Moreno et al. 2017); The study was carried out in two Nahuas communities established in different types of vegetation in the Huasteca region in the state of Hidalgo, Mexico.

MATERIAL AND METHODS

Study area

The Huasteca is a region located in central-eastern Mexico that includes portions of the states of Hidalgo, Puebla, Querétaro, San Luis Potosí, Veracruz and Tamaulipas. The majority of the inhabitants of the Huasteca region are mestizos, but a number of native peoples share the territory; Nahuas, Huastecos, Pames, Otomíes, Tepehuas and Totonacos (Olivier 2008).

The present study involves Nahuas, as they are one of the most numerous indigenous peoples and among the most widely distributed in Mexico. In the Huasteca, Nahuas live in more than 50 municipalities (county equivalents) in the states of Hidalgo, San Luis Potosí, and Veracruz (INALI 2008). In the state of Hidalgo, they live in 13 municipalities in the northeast as well as a small group in the southwest (Báez et al. 2012). The Huasteca part of Hidalgo state mainly contains two types of vegetation; tropical semi-evergreen forest (TSEF) and montane cloud forest (MCF), the TSEF covering more area than the MCF (Puig 1991). In addition, there are extensive areas of farmland, both crop and livestock. The main crops are corn, coffee, beans, sugar cane and oranges (Barthas 1996).

The study areas were selected using the criteria 1) ethnic identity, 2) more than 80% indigenous language speakers, and 3) low disturbance of the vegetative cover. This approach allows a selection of study sites associated conditions of research interest and not a priori selection assigned by the investigator (Gutiérrez-Santillán et al. 2019a). Two communities were selected: a) El Barco, in the municipality of Lolotla (21°10'45.33" – 21°10'32.89" N,

98°43'26.58" – 98°43'13.90" W; 140 masl; vegetation type TSEF, annual temperature: 18°C to 25°C, annual precipitation: 1600–1900 mm; with a populations of 380 inhabitants (INEGI 2010). In this community, the main activity is agriculture, to lesser extent livestock and fishing as a recreational activity. Some informants mentioned working in the citrus groves near the community; Figure 1); b) Tlamamala, in the municipality of Huazalingo (20°58'15.27" – 20°58'0.64" N and 98°32'40.45" – 98°32'21.34", 960 masl, vegetation type MCF, annual temperature: 13°C to 21°C, annual precipitation 1500–2800 mm; with a population of 750 inhabitants (INEGI 2010; Figure 1). The main activity is family farming, coffee harvest (coffee under shade) and livestock. Some of the informants mentioned working in the nearby city of Huejutla de Reyes, Hidalgo (approximately 40 km), and even in other cities of the country.

Fieldwork

Before fieldwork commenced, the study proposal was submitted to local authorities and approved by a community assembly (International Society of Ethnobiology 2006; <http://ethnobiology.net/code-of-ethics/>). At each interview, consent was obtained from the informant, including consent for photography and participant observation (Albuquerque et al. 2014). The field research was carried out during a two-year period from January 2011 to December 2012, with a total of 20 visits to each of the two locations.

Ethnobiological data were obtained by combining the ethnographic percentage method (10% of the population; Bernard 2006) and the snowball technique (Brewer 1995; 2002; Trotter et al. 2001). The combination of both methods allowed us to




Figure 1. Location of study sites. The red zone on the left represents the Huasteca portion of the state of Hidalgo, Mexico. The triangle shows the location of El Barco (TSEF = tropical semi-evergreen forest, 140 masl) and the star shows the location of Tlamamala (MCF = montane cloud forest, 960 masl).

work with an ethnographic sample determined by chance, having informants in general and not only expert informants. During the development of the interviews in many of the cases, the same informant was interviewed several times. Also, it should be noted that no additional ethnographic data were obtained; only sex, age and yes, his mother tongue was Nahuatl.

A total, 125 interview combinations were applied; a) 50 informants in the town of El Barco of whom 50% were women and 50% men, with age between 21 and 72 years and 94% are speakers of the indigenous language. B) 75 informants in the locality of Tlamamala of which 38% were women and

62% men, with ages between 20 and 84 years, and 88% are speakers of the indigenous language.

Ethnographic tools were also used by combining: multiple free listings (Brewer 1995; 2002) and semi-structured interviews (Bernard 2006; Albuquerque et al. 2014). First, informants were asked about plants, fungi, and animals are known in the area (multiple free listing in Spanish and Nahuatl); Later using the semi-structured interview, the use of the referred ethnospecies was asked. The interview data organized in databases, for a) plants, b) and c) fungi) animals, is organized by last taxonomic group (fish, amphibians/reptiles, birds and mammals).

Each database consisted of several sets of information: a) information the informant (name, gender, age, indigenous language speakers), b) taxonomic data (order, family, genus, species), c) nomenclature traditional (name in Nahuatl and Spanish) and d) category of use.

Collection, processing and identification of ethnobiological material

The biological material was processed and identified according to the taxonomic group in general; they had a) collections of fungi and plant specimens were collected in the field during the trips with the inhabitants to their daily activities (participative observation; Bernard 2006). The collections were processed and dehydrated adequately, later in the laboratory they were identified with the use of taxonomic keys. Fungal specimens are deposited in the collection ethnomycology "Dr. Teófilo Herrera Suárez" at the Center for Biological Research, the Autonomous University of the State of Hidalgo, Mexico. The deposit of the plants was not possible in an official collection because you do not have an herbarium of ethnobotany.

For different groups of animals fieldwork was based primarily on: b) animals or parts of animals donated by the informants; c) photographs of animals or parts of animals; d) visual stimuli – pictures based on the design of posters of species already reported in the region (Bernard 2006; Albuquerque et al. 2014), and e) field guides, e.g., birding guides (Albuquerque et al. 2014). In general, for all taxonomic groups, it has been made f) traditional nomenclature associated (documented by us) with bibliographic ethnobiological information for the region; and g) the

association of the species (identified by us) concerning published regional taxonomic lists.

Species were identified to the lowest possible taxonomic level, based on the respective literature for the group; fungi (Bautista-Nava 2007; Isidoro-Reséndiz 2011), plants (Andrade-Cetto 2009; Luna et al. 1994; Pennington and Sarukhán 2005; Pérez-Escandón et al. 2003; Puig 1991; Villavicencio 2005; Villaseñor 2016), crustaceans (Álvarez et al. 2012), mollusks (Correa-Sandoval 2003), millipedes (Bueno 2012) and vertebrates (fish: González et al. 2010; Miller 2009; amphibians and reptiles: Ramírez-Bautista et al. 2014; birds: Howell and Webb 1995; Peterson and Chalif 2000; Martínez-Morales 2007; Martínez-Morales et al. 2007; and mammals: Ceballos and Oliva 2005).

In addition, lists of species reported for the respective taxonomic groups in the Huasteca Hidalguense were used (Ramírez-Bautista et al. 2017). In the MCF, 336 species of plants have been recorded (Luna et al. 1994), 181 bird species (Martínez-Morales 2007), 34 mammal species (Mejenes-López 2008; Mejenes-López et al. 2010), 73 amphibian and reptile species (Ramírez-Bautista et al. 2014), and 22 species of fish (González-Rodríguez et al. 2010). For the TSEF, 274 species of plants have been recorded (Puig 1991; Villavicencio 2005; Villaseñor 2016), 173 bird species (Martínez-Morales et al. 2007), 35 mammal species (Mejenes-López 2008; Mejenes-López et al. 2010), 32 amphibian and reptile species (Ramírez-Bautista et al. 2014) and 28 species of fish (González-Rodríguez et al. 2010).

The categories of use were established through the use reported by informant (emic categories for region) additionally other proposals in the literature were consulted

(Farfán et al. 2007; Lira et al. 2009; Monroy-Vilchis et al. 2008). Eleven categories were identified: ceremonial, fuel, edible, construction, timber, pets, medicinal, ornamental, commercial/sale, agricultural use and other (which contained any other uses which were mentioned only once in the interviews).

Data analysis

The inventory of biocultural diversity was measured by means of indexes adapted from metrics used to analyze biological diversity. To determine whether the number of selected informants was sufficient to obtain a complete inventory of known and exploited diversity (frequency of mention, FM), an accumulation curve was plotted using the non-parametric first-order Chao index (Chao1), calculated by means of the Estimates 8.0 program (Corwell et al. 2004). The Chao1 estimator is a method that uses abundance data and is based on the number of species that occur only once or in one sample and the number of species that occur exactly twice or in two samples (Escalante 2003).

In this case, the abundance data of the species in a sample was replaced by FM data; that is, the number of times a species i is mentioned in a sample (Pineda and Verdú 2013). An approximation of the expected number of ethnospecies was calculated, to estimate whether the number of interviewees (sampling effort) was sufficient for a complete inventory. Nonparametric estimators have their statistical basis in techniques for estimating the number of classes from samples and capture–recapture techniques (Chao and Lee 1992; Jiménez-Valverde 2003). In the present case, this corresponds to informants–mentioned ethnospecies,

considering each interviewee as a sampling unit. An assumption of these nonparametric estimators is that the probability of capture – here the probability that an ethnospecies is mentioned – must remain constant throughout the period of ethnographic sampling.

The analysis of biocultural diversity was calculated based on the diversity formula proposed by Jost (2006), which is known as a zero-order and first-order measure of diversity (Hill 1973), using the formula:

$${}^qD = \left(\sum_{i=1}^s p_i^q \right)^{\frac{1}{1-q}}$$

where p_i is the abundance of species i divided by the total sum of the abundances of the S (species), and the exponent q is the order of diversity. The order of diversity (q) is influenced by the relative abundance of the species in the index; that is, the predominance of common species or of rare species. In the present case, the relative abundance values in the ecological inventories are replaced by the values obtained for the FM of the reported ethnospecies.

Zero-order (0D) and first-order (1D) diversity values were calculated. Zero-order diversity (0D) is based on the number of incidences of the species in the sample, which is equivalent to species richness, or in our case ethnospecies richness. In first-order diversity (1D), all the species in a sample are aggregated with a value exactly proportional to their relative abundance, without overvaluing the rare or common species (Hill 1973; Jost 2006; Moreno et al. 2011; Moreno et al. 2017). First-order diversity (1D) can be interpreted as the number of effective species, these being understood as units corresponding to the

numbers of species with the abundances that would theoretically coexist in a community with maximum equality (Moreno et al. 2017). Using this index facilitates comparing the numbers of species between communities and quantifying the differences between them (Hill 1973; Jost 2006; Moreno et al. 2011).

To compare the sites in terms of ethnospecies composition, the Sørensen similarity index (I_s) was used. The degree of association between the species recorded in the literature for the region (independent variable) and the ethnospecies reported in this study (response variable) was tested by linear correlation using the Past 3.20 program (Hammer et al. 2001).

In order to describe the association between use categories and biological groups, a correspondence analysis (CA) was carried out in the Statistica program (StatSoft 2004). A data matrix was created, where the variables correspond to biological groups and the cases to use categories. The chi-square statistic (X^2) and the percentage of variation explained between the first and second dimensions were calculated.

Table 1. Species–ethnospecies data for biological groups listed by inhabitants of two Nahua communities in the state of Hidalgo, Mexico. There was a strong relationship overall between the species and their cultural counterparts (ethnospecies); however, in the case of invertebrates, taxonomic identification was only made to the level of gender and family. TSEF = tropical semi-evergreen, MCF = montane cloud forest.

Biological group	Number of families	Total number of species	Number of ethnospecies TSEF	Number of ethnospecies MCF
Invertebrates	2	0	2	2
Fish	10	31	24	8
Amphibians & Reptiles	10	21	22	13
Birds	27	61	57	44
Mammals	17	34	31	34
Fungi	7	13	12	8
Plants	87	223	207	199
TOTAL	160	383	355	308

RESULTS

Analysis of Biocultural Diversity

A total of 408 ethnospecies that integrate the biocultural diversity recognized by the Nahuas of the Huasteca region in the state of Hidalgo were recorded. These correspond to 383 species distributed in 343 genera belonging to 160 families (Supplementary Material). The biological groups recorded overall were plants (59%), birds (16%), mammals (9%), fish (8%), amphibians and reptiles (5%), fungi (3%), and invertebrates (molluscs, crustaceans and millipedes >1%). The biological group with the most recorded taxonomic families was plants (58%), followed by birds (16%) and mammals (9%) (Table 1). In addition, 342 ethnospecies were mentioned in some use category; 304 species used in the TSEF and 290 species used in the MCF, a difference of 14.

The ethnospecies accumulation curves for the TSEF showed that 82% of the expected ethnospecies were represented (Chao1 = 291), and for the MCF, 79% were

represented (Chao1 = 275). In both cases, the actual ethnographic sampling effort (number of interviews) is located in the asymptote of the curve, which suggests that the inventory of known and used biodiversity was very few interviews short of a complete inventory (Figure 2).

For the TSEF, zero-order diversity (0D) was 355 ethnospecies (invertebrates 2, fish 24, amphibians and reptiles 22, birds 57, mammals 31, fungi 12 and plants 207) and first-order diversity (1D) was 234.2 ethnospecies (invertebrates 1.9, fish 21.5, amphibians and reptiles 19.9, birds 44.7, mammals 23.7, fungi 9.9 and plants 136.7). For the MCF, the 0D was 308 ethnospecies (invertebrates 2, fish 8, amphibians and reptiles 13, birds 44, mammals 34, fungi 8 and plants 199) and 1D was 202.3 ethnospecies (invertebrates 1.8, fish 6, amphibians and reptiles 8.6, birds 31, mammals 23.5, fungi 5.8 and plants 131). The observed difference between the TSEF and the MCF was 47 ethnospecies for 0D and 31.95 ethnospecies for 1D (Figure 3).

We found a similar composition of known and used ethnospecies in the two Nahua communities even though they are in

different vegetation types. The Sørensen similarity between the TSEF and the MCF was 84%, indicating that a large percentage of ethnospecies are shared between the two sites. In addition, a positive correlation was found between the number of species reported in the literature for the different taxonomic groups and the number of ethnospecies documented in the present study ($r= 0.8424$, $p= 0.0037$). This indicates that most species have their cultural counterpart and that much of the knowledge about biodiversity is shared, independently of vegetation type, a phenomenon associated with the cultural origin of the human group and the fact that both communities are in the same geographical region.

Use Trends

A total of 11 use categories were identified for both communities. The categories containing the highest number of ethnospecies are commercial trade or sale (28.7%), edible uses (17.5%) and medicinal uses (14.1%); followed by the categories with moderate values: ornamental (11.1%),




Figure 2. Accumulation ethnospecies curve based on the number of interviews and the non-parametric Chao first-order accumulation function (Chao1). On the x-axis, the number of informants is shown by community, while the y-axis shows the cumulative number of mentioned species (FM). The bars at each point represent the 95% confidence interval. TSEF = tropical semi-evergreen, MCF = montane cloud forest.




Figure 3. Diversity values of order zero (0D) and first order (1D) by vegetation type. To graph the values, the Shannon index was used in both levels of diversity. The data were the number of ethnospecies by biological group for zero-order diversity and the inverse of the Shannon index for first-order diversity. This procedure enabled the confidence intervals to be obtained. TSEF = tropical semi-evergreen, MCF = montane cloud forest.

fuel (6.3%) and pets (5.3%). The categories with the lowest values are ceremonial and construction with 4.5% each, timber (2.0%) and agricultural use (1.8%). Four of the categories (fuel, construction, timber, agricultural use) only contained plant species, and the pet category was exclusive to animals, mainly birds and mammals (Table 2).

The correspondence analysis (CA) showed the association between biological groups and anthropocentric categories, including ethnospecies that are used in different ways in the two communities. There is a significant relationship between biological group and use category ($\chi^2=537.35$, $df=50$, $p < 0.05$). The two dimensions explain 81% of the variation in the data. In the first dimension, plants contribute the most to the variance (eigenvalue= -0.4621), followed by birds (eigenvalue= 0.2648) and mammals (eigenvalue= 0.1520). In the second dimension, the main contributions to the variance are from fish (eigenvalue= -0.1816), fungi (eigenvalue= -0.0570) and amphibians and reptiles (eigenvalue= 0.0330).

We can observe in the graph (Figure 4) that certain categories of use are exclusive of certain biological groups. For example, the categories of agricultural uses, fuel, construction, and wood are exclusive to the plants; while the pet category with birds, mammals and reptiles. In turn, birds and mammals are more related to ornamental use and as pets, while reptiles with ceremonial uses. Some generic categories are more associated with vertebrates than with plants (e.g., ornamental and ceremonial), or, for example, the edible category (which should be generic) has a more significant relationship for fungi and fish; this is due to the fact that the ethnospecies that form these two biological groups generally only fulfill alimentary functions (Figure 4). The association of the biological groups and the categories of use, in this case, is given by the frequency of mention of each one of the ethnospecies for each one of the categories; so, the relationship shown in the chart (Figure 4) is specific for these two study communities.

Table 2. Relationship between number of ethnospecies mentioned and categories of use. Plants are the group with greatest diversity of categories, invertebrates and fish are the group with the least.

Biological group	I Construction	II Timber	III Agriculture	IV Fuel	trade	VI Others	Ceremonial	Ornamental	IX Medicinal	X Pets	XI Edible
Invertebrates									2		3
Fish								2	11		31
Amphibians & Reptiles						8	4		7	3	2
Birds					25	1	17	37	32	29	16
Mammals					20	4	4	27	20	17	21
Fungi					6						11
Plants	42	19	17	58	236	21	17	36	58		77
Total	42	19	17	58	287	34	42	102	130	49	161

Legend: Relationship between number of ethnospecies mentioned and categories of use. Plants are the group with greatest diversity of categories, invertebrates and fish are the group with the least.




Figure 4. The correspondence analysis shows the association between the biological groups; mammals, birds, reptiles, fish, fungi and plants, with respect to the use categories: I= construction, II= timber, III= agricultural use, IV= fuel, V= sale or commercial trade, VI= others, VII= ceremonial, VIII= ornamental, IX= medicinal, X= pets, and XI= edible. In some cases, a specific relationship is observed between biological groups and certain categories; for example, fish and fungi are associated with the edible category because fungi are normally only harvested for food, and fish can only be sold sometimes, so their main use is as food. In contrast, mammals, birds and reptiles are mostly related to ceremonial, medicinal, ornamental and pet uses, while plants are associated with uses such as timber, agriculture, construction, fuel and commercial use.

DISCUSSION

Biocultural diversity of the Nahua

The Nahuas of the Huasteca region in Hidalgo state have extensive knowledge about the biodiversity in their environment. In this study, 408 ethnospecies were reported. They are associated with 383 species belonging to a variety of biological groups (invertebrates, amphibians and reptiles, fish, birds, mammals, fungi and plants). This is a higher number of ethnospecies than were found by other comparable ethnobiological studies (Caballero and Mapes 1985; Farfán et al. 2007; Nabhan et al. 1982). Other studies have generally not addressed all three main ethnobiological subdisciplines; ethnomycology, ethnobotany and ethnozoology. Our research is a contribution from a comprehensive ethnobiology perspective, combining the three subdisciplines integrating various biological groups and evaluating social-ecological biodiversity knowledge held by a specific indigenous group. In addition, this study contributes to the analysis of biocultural diversity by treating ethnospecies as units of analysis, as has been done in ecology (Moreno 2011; Moreno et al. 2017) and shows how metrics used in the analysis of biological diversity can be adapted to the analysis of biocultural diversity.

Comprehensive ethnobiology research has been previously addressed by Toledo et al. (1983) who documented 410 species for the Pu'rhépechas, Hunn (2008) who recorded a total of 1,379 species (fungi, plants, invertebrates and vertebrates) known and used by the Zapotecs, and Aldasoro-Maya (2012) who reported 264 species known and used by the the Tlahuica. However, there are few such ethnobiology studies using a comprehensive approach, for

reasons such as lack of availability of time, funding, access to communities, or lack of transdisciplinarity, among others. It is more common to find multitaxonomic research addressing two or three biological groups (Nabhan et al. 1982; Caballero and Mapes 1985; Farfán et al. 2007).

On the other hand, regarding the knowledge and use of biocultural diversity, a high similarity is determined even though the communities are established in two different types of vegetation (TSEF and MCF). The similitude may be given because they share common species, which are generally widely distributed or introduced. It also suggests that this phenomenon is associated with the cultural memory or bio-cultural patrimony of both communities, as it belongs to the same indigenous group (Olivier 2008), as well as geographic proximity (see map). So, when studying two types of dominant vegetation in the Huasteca region (Puig 1999), it is likely that the social-ecological relationships for the Nahuas are similar.

The biocultural diversity reported here indicates a high degree of recognition and use of the natural environment in communities where the traditional language and many of the customs are still preserved. This is confirmed by the significant positive correlation between the number of ethnospecies and the number of species reported in the literature for the various taxonomic groups (Ramírez-Bautista et al. 2017).

Conceptual contribution

We propose that Biocultural diversity as: “the variety of organisms that are known, named, classified, organized, used, exploited, domesticated and/or manipulated by different human societies; including the social-ecological systems of which this

diversity forms a part at various spatiotemporal scales (Gutiérrez-Santillán 2018). But this concept also must be referred to the correlation between its biological, ecological, environmental, geographical, cultural and linguistic components; include a regional or global spatial scale (Maffi 2005; 2007; Loh and Harmon 2005; Harmon and Loh 2010; Stepp et al. 2004; 2005; Turvey and Pettoelli 2014). However, in *sensu stricto* has been seen that biological diversity and biocultural diversity are not entirely congruent in terms of the factors that generate them (Cardillo et al. 2015).

However, the design of a conceptual framework in the metric of cultural diversity analogous to that of biological diversity will not only allow us to understand it, but also to confirm that its identification and evaluation are adequate (Maclaurin and Sterelny 2008). Currently, the existence of biocultural diversity as a real and inherent property of the social-ecological relationship is not discussed, however, semantic, conceptual and analytical gaps are detected; for which different approaches can be useful to generate new research perspectives as in ecology (Moreno et al. 2011).

One of the fundamental bases for the conceptual development of biocultural diversity is the establishment of real and identifiable social-ecological units or entities, as for example in ecology, that operational taxonomic units have been used (Krell 2004). Although ethnobiology has sought to develop research on the knowledge and use of biodiversity, some of the previous studies do not discriminate between species and ethnospecies, or do not consider this criterion as a rule in ethnobiological research; taking both as an independent record or identifiable entity. We consider it essential to establish the units of analysis,

referring to the ethnospecies, which must be integrated by the taxonomic identity of the species plus their corresponding culture; this criterion should apply regardless whether the investigation has a qualitative or quantitative approach. In addition, the establishment of ethnospecies as units of analysis, as has been done in ecology (Moreno et al. 2011; Moreno et al. 2017) favors the adaptation of current metrics used in the analysis of biological diversity to the analysis of biocultural diversity.

The identity of the social-ecological units is critical to begin to understand biocultural diversity from a more local and analogous view to biodiversity. Our study allows us to observe another pattern more like those of biological diversity, such as the recognition of common species, as well as rare species (Turner et al. 2011). As examples of common species we have the coyote/coyochichi (*Canis latrans*), the pigeons/singuilotl (*Claravis pretiosa*, *Leptotila verreauxi*, *Geotrygon albifacies*), the fungus orejita-de-vejita/cuapetachiquinte (*Auricularia auricula*, *A. delicata*, *A. fuscossuccinea*), and trees such as guava (*Psidium guajava*) and framboyán/ framboyánxuchitl (*Delonix regia*). And as rare species we have the anguila/coatlmichi (*Anguilla rostrata*), the jaguar/tecuaní (*Panthera onca*) and, the perro-de-agua/atlchich (*Lontra longicaudis*), these recently reported species for the state of Hidalgo (Aguilar-López et al. 2015; Morales-García and Acosta-Rosales 2015). Or tree species that are very scarce in the surrounding forests, but culturally highly valued as the palo-escrito (*Dalbergia palo-escrito*) or the palo-varón/tlacacuahuitl (*Ulmus mexicana*). Even the documentation of a locally extinct species was obtained: the mono/chango (*Ateles geoffroyi*), which is estimated to have been extirpated from northeastern Mexico (Ceballos and Oliva

2005).

Biocultural diversity is part of the memory or biocultural heritage of indigenous peoples, their documentation, analysis and evaluation; it can be appreciated by biologists of conservation, by documenting traditional knowledge and practices historical and currently conducted on biodiversity, helping to understand changes in local biodiversity and designing appropriate conservation strategies (Brook and McLachlan 2008, Saslis-Lagoudakis and Clarke 2012). In addition, under the perspectives projected in this research, we seek to open a way for conceptualization and the establishment of methodologies that help to understand biocultural diversity from the low level to more complex analyzes, such as, for example, macro-ethnobiological (Gutiérrez-Santillán et al. 2019b), the evolutionary ethnobiology (Albuquerque y Ferreira-Junior 2017), the niche construction theory and ethnobiology (Albuquerque et al. 2018); and the development of the social-ecological theory of maximization (Albuquerque et al. 2019).

The use of new metrics for the analysis and evaluation of biocultural diversity

The comprehensive analysis of biocultural diversity (multitaxonomic and quantitative) generates a clearer picture of social-ecological knowledge. To do so, it is essential to evaluate biocultural diversity using metrics, which give greater robustness to the results and allow for comparisons, as has been done in ecology (Moreno et al. 2017).

It is, of course, important that the theoretical assumptions required for the chosen metrics be met. In this case, we evaluate whether the ethnographic sampling

effort is enough using accumulation curves (Alves et al. 2016). The curves showed that the number of interviews was enough for the inventory. This suggests that when the percentage ethnographic method, which generally implies sampling 10% of the inhabitants of a population (Bernard 2006) is applied in combination with the snowball method (Trotter et al. 2001), it is possible to obtain a robust inventory of ethnobiological information. In addition, selecting informants using the snowball method enables a random sample to be drawn (Trotter et al. 2001). In addition, the combination of ethnographic tools (multiple free lists and semi-structured interviews) also allow an adequate collection of information.

These metrics can be used to corroborate the completeness of the ethnobiological data, since the accumulation curves and estimates of chosen species (non-parametric) are good indicators of the quality of the sample (Albino-García et al. 2011; Benz et al. 2000; Hopkins and Stepp 2012; Lozada et al. 2006; Pineda and Verdú 2013), in our case indicating proper ethnographic sampling.

In the accumulation curves, frequency of mention is used, considering each informant as a sampling unit, and the mentions of ethnospecies represent their cultural importance, not their abundance in nature. In ethnographic sampling there are ethnospecies that are constantly mentioned, or common, and ethnospecies that are mentioned only occasionally or rarely. However, it is important to assess in which cases to apply the accumulation curve or not, since the curve may be indicating that our inventory is close to the optimum; that is, increasing the sampling number maintains the asymptote constant (Alves et al. 2016; Moroy-Vilchis et al. 2008); but other methodological aspects must be adhered to,

such as ensuring random and homogeneous sampling.

For the implementation of these and other metrics, it is essential to keep in mind that the methods used in the analysis of biological diversity are constantly being updated (Moreno et al. 2011; 2017), making it necessary to evaluate their usefulness in ethnobiological or biocultural diversity studies. For example, the zero-order (0D) and first-order diversity index (1D) are in current use (Hill 1973; Jost 2006). The advantage of using these indices is that the results are comparable within a study and with other studies in the same region whether intra- or inter-ethnically; and across time spans to evaluate cultural changes in knowledge or uses; that is, to evaluate local knowledge at different space and time scales (Gutiérrez-Santillán et al. 2019b).

In this study we use only the 0D and 1D diversity indices, because the zero-order diversity data corresponds to species richness, in our case ethnospecies richness, while first-order diversity includes all species with a weight precisely proportional to their abundance in the community (Hill 1973; Jost 2006; Moreno et al. 2011). In ethnographic sampling, all species reported by the inhabitants correspond to a site, and they have the same representation in the sample regardless of their frequency of mention, so this proportion is maintained. For this index, second-order diversity (2D) is also calculated, but its application is not recommended for the present case, because this measure is based on the dominance of the species present in a sample (Jost 2006; Moreno et al. 2011; 2017). In ethnobiological data we have ethnospecies that are represented with a high frequency of mention, which may be biased towards some cultural or even economic preference that strongly influences its frequent mention,

which we consider could cause bias.

We suggest that the suitability of biodiversity metrics should be evaluated for use in ethnobiological studies, in order to determine whether an ethnographic sample behaves in a similar way to an ecological sample. The relevance of other indexes should be considered as well; not only those that are based on incidence but also those that take abundance values into account. We consider that using ethnobiological or social-ecological units (taxonomic species + cultural assignment) and making comparisons between the degree of biological diversity and the diversity of ethnospecies are sufficient reason to apply more refined forms of analysis to evaluate biocultural diversity. However, it is essential to ensure the quality of the methodological design such that it involves proper collection of sufficient data, to avoid a subjective interpretation.

Trends of use

One of the most important characteristics of biocultural diversity is its use by human groups. Studies in other regions of Mexico have documented various use categories for different taxonomic groups (Lira et al. 2009; Monroy-Vilchis et al. 2008). In this study, 11 categories, which contribute to satisfying a wide range of human needs, were identified. Different indigenous groups use species in different ways for different purposes, which reflects the importance of nature in contemporary societies (Alonso-Castro et al. 2011; Carbajal-Esquivel et al. 2012; Toledo 1994).

In addition to the application of appropriate metrics, the use of multivariate statistical methods provides a broader view of the social-ecological relationships in the study area. In this case, implementing CA

enabled a graphical association to be made between use categories and the various taxonomic groups. Several authors have used these methods, finding structural relationships between the study groups; e.g., between taxonomic families and habitat types (Molares and Ladio 2009), species management types (Parra et al. 2012), species (Zamudio and Hilgert 2012) and knowledge among several indigenous groups (Núñez-García et al. 2012). We found that certain biological groups are associated with specific categories, as a characteristic of the biological group and its use.

CONCLUSIONS

The Nahuas of the Huasteca region in the state of Hidalgo have extensive knowledge and make widespread use of the biodiversity in their territories. This study has shown that each of the vegetation types contributes particular characteristics in terms of social-ecological relationships at the cultural level, a phenomenon associated with the presence of unique species of the tropical semi-evergreen forest and the montane cloud forest. However, since the two study communities belong to the same indigenous group, are located in the same geographical region and are in contact with the same widely distributed and introduced species, there are strong similarities in their knowledge and use of these species. In addition, it was found that each of the biological groups contributes in a particular way and in some cases exclusively in their respective use categories, especially in the case of plants and animals.

This study contributes to a comprehensive view of the multitaxonomic analysis of biodiversity known and used by a indigenous group, by combining the three

main ethnobiological subdisciplines: ethnobotany, ethnomycology and ethnozoology (Figure 5). It is suggested that the conceptualization of biocultural diversity be extended beyond its cultural and biological components. We propose that biocultural diversity be considered as “the variety of organisms that are known, named, classified, organized, used, exploited, domesticated and/or manipulated by different human societies”; including the social-ecological systems of which this diversity forms a part at various spatiotemporal scales (Gutiérrez-Santillán 2019b).

It is composed essentially of ethnospecies, which consist of a taxonomic entity (species) together with its cultural counterpart (traditional name). When the basic units of biocultural diversity are defined, they can be analyzed using the new metrics used for the analysis of biological diversity, enabling a better assessment of biocultural diversity and more robust data to be generated, which can be compared on the intra- or interethnic level, between regions, ecosystems, or other divisions. In addition, this approach can be used to generate and provide sociocultural data for the implementation of conservation strategies under a biocultural approach.

ACKNOWLEDGEMENTS

Thanks are expressed to the residents of the study communities who kindly supported this research and contributed their invaluable knowledge, and to the UAEH for its support of AMF, ASG and GSR so that this research could be carried out. TVGS received a scholarship from CONACyT (389096/254575).



Figure 5. A) Pet boar (*Tayassu tajacu*, MCF); B) Striped (lowland) paca meat (*Cuniculus paca*, TSEF); C) Child with preserved margay (*Leopardus weidii*, MCF); D) Pet red-billed pigeon (*Patagioenas flavirostris*, TSEF); E) Informant with preserved emerald toucanet (*Aulacorhynchus prasinus*, MCF); F) Child fisherman (TSEF); G) crayfish (*Procambarus* sp., TSEF); H) cosol, river crayfish trap (TSEF); I) Cross decorated with cycad leaves (*Zamia herrerae*, MCF); J) Itztacnanacatl fungus on papatla leaf (*Pleurotus albidus*, *Heliconia collinsiana*; MCF); K) yam and squash (*Ipomoea batatas*, *Cucurbita maxima*; TSEF); L) two species of marigold (cempoal) in ceremony for festival of Our Lady of Guadalupe (Dec. 12) (*Tagetes coronopifolia*, *Tagetes erecta*; TSEF). TSEF = tropical semi-evergreen, MCF = montane cloud forest.

REFERENCES

- Aguilar-López M, Rojas-Martínez A, Cornejo-Latorre C, Sánchez-Hernández C, Vite-Silva VD, Ramos-Frías J (2015) **Registros notables de mamíferos terrestres del estado de Hidalgo, México.** *Acta Zoológica Mexicana* (n.s.) 31(3): 403-411.
- Albino-García C, Cervantes H, López M, Ríos-Casanova L, Lira R (2011) **Patrones de diversidad y aspectos etnobotánicos de las plantas arvenses del valle de Tehuacán-Cuicatlán: el caso de San Rafael, municipio de Cioxcatlán, Puebla.** *Revista Mexicana de Biodiversidad* 82:1005-1019.
- Albuquerque UP, Medeiros PM, Ferreira-Junior W, Silva TC, Silva RR, Goncalves-Souza T (2019) **Social-ecological theory of maximization: basic concepts and two initial models.** *Biological Theory* doi.org/10.1007/s13752-019-00316-8.
- Albuquerque UP, Santos-Gonçalves PH, Ferreira-Júnior W, Silva-Chaves L, Silva-Oliveira RC, Lima-Silva TL, DosSantos GC, Lima-Araújo E (2018) **Humans as niche constructors: Revisiting the concept of chronic anthropogenic disturbances in ecology.** *Perspectives in Ecology and Conservation* 16(1): 1–11. doi.org/10.1016/j.pecon.2017.08.006.
- Albuquerque UP, Ferreira-Júnior W (2017) **What do we study in evolutionary ethnobiology? Defining the theoretical basis for a research program.** *Evolutionary Biology* 44(2): 206–215 doi.org/10.1007/s11692-016-9398-z.
- Albuquerque UP, Ferreira-Júnior W, Santoro FR, Torres-Avilez WM, Sousa-Júnior JR (2015). **Niche construction theory and ethnobiology.** In: Albuquerque UP, Medeiros PM, Casas A (eds). *Evolutionary ethnobiology*. Springer. New York, USA. pp. 73–87.
- Albuquerque UP, Cunha L, Lucena R, Alves RRN (2014) **Methods and techniques in ethnobiology and ethnoecology.** 1ed. Springer, New York, USA doi:10.1007/978-1-4614-8636-7.
- Alcántara-Salinas G, Ellen R, Valiñas-Coalla L, Caballero J, Argueta-Villamar A (2013) **Alternative ways of representing zapotec and cuicatec folk classification of birds: a multidimensional model and its implications for culturally-informed conservation in Oaxaca, Mexico.** *Journal of Ethnobiology and Ethnomedicine* 9:81 doi:10.1186/1746-4269-9-81.
- Alcorn JB (1981a) **Huastec noncrop resource management: implications for Prehistoric rain forest management.** *Human Ecology* 9:395-417.
- Alcorn JB (1981b) **Some factors influencing botanical resource perception among the Huastec: suggestions for future ethnobotanical inquiry.** *Journal of Ethnobiology* 1:221-230.
- Alcorn JB (1983) **El te'lom huasteco: presente, pasado y future de un sistema silvicultural indígena.** *Biótica* 8:315-331.
- Alcorn JB (1984) **Huastec mayan ethnobotany.** 1ed. University of Texas Press, Austin, U.S.A.
- Aldasoro-Maya EM (2012) **Documenting y contextualizing Pjiekakjoo (Tlahuica) knowledges through a collaborative research project.** PhD Thesis, University of Washington, Washington D.C., U.S.A.
- Alonso-Castro AJ, Carranza-Álvarez C, Maldonado-Mirya JJ, Jacobo-Salcedo MR, Quezada-Rivera DA, Lorenzo-Márquez H, Figueroa-Zúñiga LA, Fernández-Galicia C, Ríos-Reyes NA, León-Rubio MA, Rodríguez-Gallegos V, Medellín-Milán P (2011) **Zootherapeutic practices in Aquismón, San Luis Potosí, México.** *Journal of Ethnopharmacology* 138(1):233-237 doi:10.1016/j.jep.2011.09.020.
- Álvarez F, Villalobos JL, Armendariz G, Hernández C (2012) **Relación biogeográfica entre cangrejos dulceacuícolas y acociles a lo largo de la zona mexicana de transición: revaluación de la hipótesis de Rodríguez (1986).** *Revista Mexicana de Biodiversidad* 83: 1073-1083.
- Álvarez MS (2002) **Cambio sociocultural y usos de recursos forestales de las selvas, en una comunidad tepehua: El caso de Pisaflores en la Huasteca Veracruzana.** Undergraduate thesis, Escuela Nacional de Antropología e Historia, México.
- Alves MM, Faria S, Alves RRN (2016) **Wild vertebrates kept as pets in the semiarid region of Brazil.** *Tropical Conservation Science* 9:354-368.
- Andrade-Cetto A (2009) **Ethnobotanical study of the medicinal plants from Tlanchinol, Hidalgo, México.** *Journal of Ethnopharmacology* 122:163-171 doi: 10.1016/j.jep.2008.12.008.
- Argueta-Villamar A (2008) **Los saberes p'urhépecha.** Universidad Michoacana de San Nicolás de Hidalgo - Universidad Nacional

- Autónoma de México - Gobierno del Estado de Michoacán - Programa de Naciones Unidas para el Medio Ambiente, D.F., México.
- Báez L, Garret G, Pérez D, Moreno B, Fierro UJ, Hernández MG (2012) **Los pueblos indígenas de Hidalgo: Atlas etnográfico**. Instituto Nacional de Antropología e Historia - Gobierno del Estado de Hidalgo - Consejo Estatal para la Cultura y las Artes de Hidalgo, D.F., México.
- Barthas B (1996) **De la selva al naranjal (transformaciones en la agricultura indígena en la Huasteca potosina)**. In: Bovin P (ed), *El campo mexicano: una modernización a marchas forzadas*. 1 ed. Centro Francés de Estudios Mexicanos y Centroamericanos (CEMCA) - Institut Francais de Recherche Scientifique pour le Développement en Coopération (ORSTOM), D.F., pp. 183-199.
- Bautista-Nava E (2007) **Taxonomía y conocimiento tradicional de Cantharellus F. (Fungi, Cantharellaceae) en el noreste del estado de Hidalgo**. Undergraduate thesis, Universidad Autónoma del Estado de Hidalgo, México.
- Benz BE, Cevallos J, Santana F, Rosales J, Graf S (2000) **Losing knowledge about plant use in the sierra de Manantlan Biosphere Reserve, Mexico**. *Economic Botany* 54:183-191.
- Berlin B (1973) **Folk systematics in relation to biological classification and nomenclature**. *Annu Rev Ecol Syst* 4:259-27.
- Berlin B, Breedlove DE, Raven PH (1973) **General principles of classification and nomenclature in folk biology**. *Am Anthropol* 75:214-242.
- Bernard HR (2006) **Research methods in anthropology, qualitative and quantitative approaches**. 4ed. Altamira Press, Lanham, New York, Toronto, Oxford.
- Boege E. 2008. **El patrimonio biocultural de pueblos indígenas de México**. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, D.F., México.
- Brewer D (1995) **Cognitive indicators of knowledge in semantic domains**. *Journal of Quantitative Anthropology* 5:107-28.
- Brewer D (2002) **Supplementary interviewing techniques to maximize output in free listing tasks**. *Field Methods* 14:108-118.
- Brook RK, McLachlan SM (2008) **Trends and prospects for local knowledge in ecological and conservation research and monitoring**. *Biodiversity and Conservation* 17:3501-3512.
- Bueno J (2012) **Diplópodos: los desconocidos formadores de suelo**. *Biodiversitas* 102:1-5.
- Bye R, Linares E (2000) **Relationships between mexican ethnobotanical diversity and indigenous peoples**. In: Minnis PE, Elisens WJ (eds) *Biodiversity and Native America*, University of Oklahoma, Oklahoma, USA, pp. 44-73.
- Caballero N, Mapes C (1985) **Gathering and subsistence patterns among the p'urhepecha indians of Mexico**. *Journal of Ethnobiology* 5:31-47.
- Cano O (1988) **Herbolaría y Etnozoología en Papantla**. Dirección General de Culturas Populares - SEP, México.
- Carbajal-Esquivel H, Fortanelli J, García-Pérez J, Reyes-Agüero JA, Yáñez-Espinosa L, Bonta M (2012). **Use value of food plants in the Xi'iuy indigenous community of Las Guapas, Rayon, San Luis, Mexico**. *Ethnobiology Letters* 3:39-55 doi.org/10.14237/ebl.3.2012.40.
- Cardillo M, Bromham L, Greenhill SJ (2015) **Links between language diversity and species richness can be confounded by spatial autocorrelation**. *Proc.R.Soc. B* 282: 20142986 doi.org/10.1098/rspb.2014.2986
- Ceballos G, Oliva G (2005) **Los mamíferos silvestres de México**. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad - Fondo de Cultura Económica, D.F., México.
- Chao A, Lee, SM (1992) **Estimating the number of classes via sample coverage**. *Journal of the American Statistical Association* 87:210-217 doi.10.2307/2290471.
- Correa-Sandoval A (2003) **Gastrópodos terrestres del Noreste de México**. *Revista de Biología Tropical* 51 (Suppl. 3):507-522.
- Corwell RK, Mao CX, Chang J (2004) **Interpolating, extrapolating, and comparing incidence based species accumulation curves**. *Ecology* 85:2717-2727.
- De Ávila A (2008) **La diversidad lingüística y el conocimiento etnobiológico**. In: *Capital Natural de México*, vol. I: Conocimiento actual de la biodiversidad. 1 ed. CONABIO, México, pp. 497-556.

- Dunn CP (2008) **Biocultural diversity should be a priority for conservation.** *Nature* 456: 315.
- Elands BHM, Vierikko K, Anderson E, Fischer Lk, Gonçalves P, Haase D, Kowarik I, Luz AC, Niemelä J, Santos-Reis M, Wiersum KF (2018) **Biocultural diversity: A novel concepts to assess human-nature interrelations, nature conservation and stewardship in cities.** *Urban Forestry & Urban Greening* doi.org/10.1016/j.ufug.2018.04.006.
- Escalante T (2003) **¿Cuántas especies hay? Los estimadores no paramétricos de Chao.** *Elementos*, 52:53-56.
- Farfán B, Casas A, Ibarra-Manríquez G, Pérez-Negrón E (2007) **Mazahua ethnobotany and subsistence in the Monarch Butterfly Biosphere Reserve, Mexico.** *Economic Botany* 61:173-191 doi.10.1663/0013-0001(2007)61[173:MEASIT]2.0.CO;2.
- Gallardo-Arias P (2004) **Los especialistas de la curación: curanderos teenek y nahuas de Aquismón.** *Anales de Antropología* 38:179-200.
- Garibay-Orijel R, Caballero J, Estrada-Torres A, Cifuentes J (2007) **Understanding cultural significance, the edible mushrooms case.** *Journal of Ethnobiology and Ethnomedicine* doi:10.1186/1746-4269-3-4.
- Gavin MC, McCarter J, Mead A, Berkes F, Stepp JR, Peterson D, Tang R (2015) **Defining biocultural approaches to conservation.** *Trends in Ecology & Evolution* 30:140-145 doi.org/10.1016/j.tree.2014.12.005.
- González KA, Ramírez A, Sánchez E, Montañón SE (2010) **Los peces de la huasteca hidalguense.** Universidad Autónoma del Estado de Hidalgo - Fondo Mixto CONACyT - Gobierno del Estado de Hidalgo, México.
- Gorenflo LJ, Romaine S, Mittermeier R, Walker-Painemilla K (2012) **Co-occurrence of linguistic and biological diversity in biodiversity hotspots and high biodiversity wilderness areas.** *Proceedings of the National Academy of Sciences of the United States of America* 109:8032-8037 doi.10.1073/pnas.1117511109.
- Grasser S, Schunko C, Vogl CR (2012) **Gathering “tea” from necessity to connectedness with nature. Local knowledge about wild plant gathering in the Biosphere Reserve Grosses Walserta (Austria).** *Journal Ethnobiology and Ethnomedicine* 8:31.
- Gutiérrez-Santillán TV, Albuquerque UP, Valenzuela-Galván D, Reyes-Zepeda F, Vázquez L-B, Mora-Olivo A, Arellano-Méndez LU (2019a) **Trends on mexican ethnozoological research, vertebrates case: a systematic review.** *Ethnobiology and Conservation* 8:1 doi:10.15451/ec2019-01-8.01-1-39.
- Gutiérrez-Santillán TV, Valenzuela-Galván D, Albuquerque UP, Reyes-Zepeda F, Arellano-Uriel LU, Mora-Olivo A, Vázquez L-B (2019b). **The spatiotemporal scale of ethnobiology: A conceptual contribution in the application of meta-analysis and the development of the macro-ethnobiological approach.** In Albuquerque UP, Paiva RF, Fernandes LV, Alves RR. *Methods and Techniques in Ethnobiology and Ethnoecology.* New York, USA doi.org/10.1007/978-1-4939-8919-5.
- Gutiérrez-Santillán TV (2018) **Meta-análisis y macro-etnobiología: nuevas perspectivas de investigación etnobiológica, el caso de la etnozología mexicana.** PhD Thesis, Universidad Autónoma de Tamaulipas, México.
- Gutiérrez-Santillán TV (2013) **Diversidad biocultural y especies bioculturales clave: un estudio etnobiológico en dos comunidades nahuas en la Huasteca Hidalguense.** MSc. dissertation, Universidad Autónoma del Estado de Hidalgo, México.
- Hammer Ø, Harper DAT, Ryan PD (2001) **PAST: Paleontological statistics software package for education and data analysis.** *Palaeontologia Electronica* 4:1-9.
- Harmon D, Loh J (2010) **The index of linguistic diversity: A new quantitative measure of trends in the status of the world’s languages.** *LD&C* 4:97–151.
- Hernández L, Bautista B (2011) **Diccionario de nombres científicos náhuatl, comunes en español y comunes en inglés de cinco grupos de vertebrados de la Huasteca hidalguense.** Undergraduate Thesis, Instituto Tecnológico de Huejutla, México.
- Hernández V (2003) **Estudio etnobotánico medicinal de dos comunidades del municipio de Atlapexco en el estado de Hidalgo. Tlatemolistli tle xiupajtli ipan ome chinantli tlen Atlapech chinankotlatilanketl ipan Hidalgo Tlatokajyotl.** Undergraduate thesis, Universidad Nacional Autónoma de México, México.

- Hill MO (1973) **Diversity and evenness: a unifying notation and its consequences.** *Ecology* 54:427-432.
- Hong SK (2013) **Biocultural diversity conservation for island and islanders: Necessity, goal and activity.** *Journal of Marine and Island Cultures*. 2: 102-106 doi.org/10.1016/j.imic.2013.11.004.
- Hopkins AL, Stepp JR (2012) **Distribution of Herbal Remedy Knowledge in Tabi, Yucatan, Mexico.** *Economic Botany* 66:249-254.
- Howell SNG, Webb S (1995) **A guide to the birds of Mexico and Northern Central America.** 1 ed. Oxford University Press, New York, U.S.A.
- Hunn E (1998) **Mixtepec Zapotec ethnobiological classification: A preliminary sketch and theoretical commentary.** *Anthropologica* 40: 35-48.
- Hunn E (2008) **A Zapotec natural history: trees, herbs, and flowers, birds, beasts, and bugs in the life of San Juan Gbëë.** 1ed. University of Arizona Press, Tucson, U.S.A.
- Hunn E (2011) **Ethnozology.** In: Anderson EN, Pearsall DM, Hunn E, Turner NJ (eds) *Ethnobiology*. Wiley-Blackwell, U.S.A. pp. 83-96.
- Huntington HP (2013) **Local knowledge as a resource for Nepal: building partnerships with scientists and conservationists.** *Conservation Science* 1:1-4 doi.org/10.3126/cs.v1i1.8577.
- INALI Instituto Nacional de Lengua Indígena (2008). **Catálogo de las lenguas indígenas nacionales: variantes lingüísticas de México con sus autodenominaciones y referencias geostadísticas.** Diario Oficial de la Federación. Gobierno de México, D.F., México.
- INEGI. 2010. **Anuario estadístico del estado de Hidalgo.** Aguascalientes, Aguascalientes, México.
- International Society of Ethnobiology, 2006, <http://ethnobiology.net/code-of-ethics/>
- Isidoro-Resendiz D (2011) **Importancia cultural de los hongos comestibles en el municipio de Tepehuacán de Guerrero, Hidalgo.** Undergraduate thesis, Universidad Autónoma del Estado de Hidalgo, México.
- Jaimes MI, Gómez G, Pacheco N, Reyes SR (2014) **Uso y manejo de la avifauna en San Miguel Tzinacapan, municipio de Cuetzalan del Progreso, Puebla, México.** In: Vásquez-Dávila MA (ed.) *Aves, personas y culturas: estudios de etno-ornitología 1.* Red de Etnoecología y Patrimonio Biocultural – CONACyT - Instituto Tecnológico del Valle de Oaxaca - Cartel Editores - Universidad Tecnológica del Chocó. Oaxaca, México, pp 243-259.
- Jiménez-Valverde A (2003) **Las curvas de acumulación de especies y la necesidad de evaluar la calidad de los inventarios biológicos.** *Revista Ibérica de Aracnología* 8:151-161.
- Jost L (2006) **Entropy and diversity.** *Oikos* 113:363-375 doi.org/10.1111/j.2006.0030-1299.14714.x.
- Krell FT (2004) **Parataxonomy in biodiversity studies-pitfalls and applicability of 'morphospecies' sorting.** *Biodiversity and Conservation* 13(4): 795–812.
- Lira R, Casas A, Rosas-López R, Paredes-Flores M, Pérez-Negrón E, Rangel-Landa S, Solís L, Torres I, Dávila, P. (2009). **Traditional knowledge and useful plant richness in the Tehuacán–Cuicatlán Valley, Mexico.** *Economic Botany* 63:271-287 doi.org/10.1007/s12231-009-9075-6.
- Loh J, Harmon D (2005) **A global index of biocultural diversity.** *Ecological Indicators* 5:321–241 doi.org/10.1016/j.ecolind.2005.02.005.
- López del Toro, P., Andresen E, Barraza L, Estrada A (2009) **Attitudes and knowledge of shade-coffee farmers towards vertebrates and their ecological functions.** *Tropical Conservation Science* 2: 299-318.
- López-Villafranco ME, Aguilar-Contreras A (2010) **Etnobotánica médica de los tepehuas de Hidalgo.** In: Moreno A, Pulido MT, Mariaca R, Valadez R, Mejía P, Gutiérrez TV (eds) *Sistemas biocognitivos tradicionales: paradigmas en la conservación biológica y el fortalecimiento cultural.* Asociación Etnobiológica Mexicana A.C. - Global Diversity Foundation - Universidad Autónoma del Estado de Hidalgo - El Colegio de la Frontera Sur - Sociedad Latinoamericana de Etnobiología. D.F. México, pp 284-291.

- Lozada M, Ladio A, Weigandt M (2006) **Cultural transmission of ethnobotanical knowledge in a rural community of northwestern Patagonia, Argentina.** *Economic Botany* 60:374-385 doi.org/10.1663/0013-0001(2006)60[374:CTOEKI]2.0.CO;2.
- Luna I, Ocegueda S, Alcántara O (1994) **Florística y notas biogeográficas del bosque mesófilo de montaña del municipio de Tlanchinol, Hidalgo, México.** *Anales del Instituto de Biología, Serie Botánica* 65:31-62.
- Luque D, Doode S (2010) **Los comcáac (seri): hacia una diversidad biocultural del Golfo de California y estado de Sonora.** *Estudios Sociales 1 (Esp.):*274-300.
- Maclaurin J., Sterelny K (2008) **What is biodiversity?.** The University of Chicago Press. Chicago, USA.
- Maffi L, Dilts O (2014). **Biocultural diversity toolkit: biocultural approaches to conservation & development.** Terralingua, Salt Spring Island, British Columbia, Canada.
- Maffi L, Woodley E (2008) **Biocultural diversity conservation: a global sourcebook.** (1ed) Earthscan. Columbia.
- Maffi L (2007) **Biocultural diversity and sustainability.** In: Pretty J, Ball A, Benton T, Guivant J, Lee D, Orr D, Pfeffer M, Ward H (eds) *Sage handbook on environment and society.* Sage Publications, London, pp 267-277.
- Maffi L (2005) **Linguistic, cultural and biological diversity.** Terralingua, Salt Spring Island, British Columbia, Canada.
- Maffi L (2001) **On biocultural diversity linking language, knowledge, and the environment.** Smithsonian Institution Press, Washington, U.S.A.
- Martínez-Morales M (2007) **Avifauna del bosque mesófilo de montaña del noreste de Hidalgo, México.** *Revista Mexicana de Biodiversidad* 78:149-162.
- Martínez-Morales M, Ortiz-Pulido R, de la Barreda B, Zuria IL, Bravo-Cadena J, Valencia-Herverth J (2007) **Hidalgo.** In: Ortiz-Pulido R, Navarro-Sigüenza A, Gómez de Silva H, Rojas-Soto O, Peterson TA (Eds.). *Avifaunas Estatales de México.* CIPAMEX, México, pp 49- 95.
- Mejenes-López SMA, Hernández-Bautista M, Barragán-Torres J, Pacheco J (2010) **Los mamíferos del estado de Hidalgo, México.** *Therya* 1:161-188.
- Mejenes-López SMA (2008) **Mastofauna de la región Huasteca y Sierra de Hidalgo.** Undergraduate thesis, Universidad Nacional Autónoma de México, México.
- Miller RR (2009) **Peces dulceacuícolas de México.** CONABIO - Sociedad Ictiológica Mexicana A.C. - El Colegio de la Frontera Sur - Consejo de los Peces del Desierto México-Estados Unidos. D.F., México.
- Molares S, Ladio A (2009) **The usefulness of edible and medicinal Fabaceae in Argentina and Chilean Patagonia: Environmental availability and other sources of supply.** *Evidence-Based Complementary and Alternative Medicine* 1:12 doi:10.1155/2012/901918.
- Monroy-Vilchis O, Cabrera L, Suárez P, Zarco-González MM, Rodríguez-Soto C, Urios V (2008) **Uso tradicional de vertebrados silvestres en la Sierra Nanchititla, México.** *Interciencia* 33:308-313.
- Montaño SE, Moreno A, González KA (2010) **Revalorando la pesca náhuatl como ejemplo de una práctica sustentable: estudio de caso en la región hidalguense, México Central.** In: Moreno A, Pulido MT, Mariaca R, Valadez R, Mejía P, Gutiérrez TV (eds) *Sistemas biocognitivos tradicionales: paradigmas en la conservación biológica y el fortalecimiento cultural.* Asociación Etnobiológica Mexicana A.C. - Global Diversity Foundation - Universidad Autónoma del Estado de Hidalgo - El Colegio de la Frontera Sur - Sociedad Latinoamericana de Etnobiología. D.F., México, pp 317-321.
- Montoya JJ (1968) **Magia y cacería entre los nahuas de la Sierra de Hidalgo.** *Boletín INAH* 34:22-23.
- Morales-García AD, Acosta-Rosales A (2015) **Registros recientes de jaguar (*Panthera onca*) en el estado de Hidalgo, México.** *Revista Mexicana de Mastozoología Nueva Época* 5(2): 66-72.
- Moreno CE, Calderón-Patrón JM, Arroyo-Rodríguez V, Barragán F, Escobar F, Gómez-Ortiz Y, Martín-Regalado N, Martínez-Falcón AP, Martínez-Morales MA, Mendoza E, Ortega-Martínez IJ, Pérez-Hernández CX, Pineda E, Pineda-López R, Rios-Díaz CL, Rodríguez P, Rosas F, Schondube JE, Zuria I (2017) **Measuring biodiversity in the Anthropocene: a simple guide to helpful methods.** *Biodiversity and Conservation* 20(12): 2993-2998 doi.10.1007/s10531-017-1401-1.

- Moreno CE, Barragán F, Pineda E, Pavón NP (2011) **Reanalizando la diversidad alfa: alternativas para interpretar y comparar información sobre comunidades ecológicas.** *Revista Mexicana de Biodiversidad* 82:1249-1261.
- Nabhan GP, Rea AM, Reichhardt KL, Mellink E, Hutchinson CF (1982) **Papago influence on habitat and biotic diversity: Quitovac oasis ethnoecology.** *Journal Ethnobiology* 2: 124-143.
- Neyra-González, L. Durand-Smith L. (1998) **Biodiversidad.** In: La diversidad biológica de México: Estudio de país. Comisión para el Conocimiento y uso de la Biodiversidad, México, pp. 61-102.
- Núñez-García RM, Fuente-Carrasco ME, Venegas-Barrera CS (2012) **La avifauna en la memoria biocultural de la juventud indígena en la Sierra Juárez de Oaxaca, México.** *Universidad y Ciencia* 28(3): 201-216.
- Olivier G (2008) **Viaje a la Huasteca con Guy Stresser-Péan.** Fondo de Cultura Económica - CEMCA, D.F., México.
- Parra F, Blancas JJ, Casas A (2012) **Landscape management and domestication of *Stenocerus pruinosus* (Cactaceae) in the Tehuacán Valley: human guided selection and gene flow.** *Journal Ethnobiology and Ethnomedicine* 8:1-32 doi. 10.1186/1746-4269-8-32.
- Penguilly MA, Moreno A, Goyenechea I, Espinosa G (2010) **Percepción acerca de las lagartijas consideradas nocivas por algunos otomíes, nahuas, tepehuas y mestizos en el estado de Hidalgo, México.** In: Moreno A, Pulido MT, Mariaca R, Valadez R, Mejía P, Gutiérrez TV (eds) *Sistemas biocognitivos tradicionales: paradigmas en la conservación biológica y el fortalecimiento cultural.* Asociación Etnobiológica Mexicana A.C. - Global Diversity Foundation - Universidad Autónoma del Estado de Hidalgo - El Colegio de la Frontera Sur - Sociedad Latinoamericana de Etnobiología, D.F., México, pp 99-105.
- Pennington TD, Sarukhán J (2005) **Árboles tropicales de México, manual para la identificación de las principales especies.** Universidad Nacional Autónoma de México - Fondo de Cultura Económica, D.F., México.
- Pérez-Escandón BE, Villavicencio-Nieto MA, Ramírez-Aguirre A (2003) **Lista de plantas útiles del estado de Hidalgo.** Universidad Autónoma del Estado de Hidalgo, México.
- Peterson RT, Chalif EL (2000) **Aves de México. Guía de Campo.** Editorial Diana, D.F., México.
- Pineda R, Verdú JR (2013) **Cuaderno de prácticas. Medición de la biodiversidad: diversidades alfa, beta y gamma.** Universidad Autónoma de Querétaro - Universidad de Alicante, México.
- Piotrowska Z (2013a) **No sé si los aires ven con coraje que voy a hacer curaciones. Misión del curandero, sus desafíos y peligros según los nahuas de la Huasteca hidalguense.** In: La terca realidad. La Huasteca como espejo cultural. CIESAS - El Colegio de San Luis S.C., México.
- Piotrowska Z (2013b) **Cerros y curanderos entre los nahuas de la Huasteca.** In: La terca realidad. La Huasteca como espejo cultural. CIESAS - El Colegio de San Luis S.C., México.
- Pretty J, Adams B, Berkes F, Ferreira S, Dudley N, Hunn E (2009). **The intersections of biological diversity and cultural diversity: Towards integration.** *Conservation and Society* 7(2): 100-112 doi:10.4103/0972-4923.58642.
- Puig H (1991) **Vegetación de la Huasteca (México) Estudio fitogeográfico y ecológico.** Institut Francais de Recherche Scientifique pour le Développement en Coopération (ORSTOM) - Instituto de Ecología A.C. - Centre D'Estudes Mexicaines et Centraméricaines (CEMCA), D.F., México.
- Ramírez-Bautista A, Sánchez-González A, Sánchez-Rojas G, Cuevas-Cardona C. (2017) **Biodiversidad del Estado de Hidalgo.** 1ed Universidad Autónoma del Estado de Hidalgo, Consejo Nacional de Ciencia y Tecnología, México.
- Ramírez-Bautista A, Hernández-Salinas U, Cruz-Elizalde R, Berriozabal-Islas C, Lara-Tuñño D, Goyenechea I, Castillo-Cerón JM (2014) **Los anfibios y reptiles de Hidalgo, México: Diversidad, biogeografía y conservación.** Sociedad Herpetológica Mexicana, México.
- Ruvalcaba J, Pérez JM, Herrera O (2004) **La Huasteca, un recorrido por su diversidad.** CIESAS - El Colegio de San Luis - El Colegio de Tamaulipas, México.

- Saslis-Lagoudakis CH, Clarke AC (2013) **Ethnobiology: the missing link in ecology and evolution.** *Trends in Ecology & Evolution* 28:67–68 doi: 10.1016/j.tree.2012.10.017.
- StatSoft Inc (2004) **STATISTICA (Data Analysis Software System)** versión 7. www.statsoft.com
- Stepp JR, Castaneda H, Cervone S (2005) **Mountains and biocultural diversity.** *Mountain Research and Development* 25(3): 223–227 doi.org/10.1659/0276-4741(2005)025[0223:MABD]2.0.CO;2.
- Stepp JR, Cervone S, Castaneda H, Lasseter A, Stocks G, Gichon Y (2004) **Development of a GIS for global biocultural diversity.** *Policy Matters* 13: 267-271.
- Sutherland WJ (2003) **Parallel extinction risk and global distribution of languages and species.** *Nature* 423: 276-279.
- Toledo VM (2010) **La biodiversidad de México. Inventarios, manejos, usos, informática, conservación e importancia cultural.** FCE, CONACULTA. D.F., México.
- Toledo VM (1994) **Biodiversity and Cultural Diversity in Mexico.** *Different Drummer* 1:16-19.
- Toledo VM, Caballero J, Mapes C, Barrera N, Argueta A, Nuñez MA (1983) **Los purépechas de Patzcuaro: Una aproximación ecológica.** 2ed Serie Etnociencia: Cuadernos de Etnobiología 1 – Secretaría de Educación Pública - Dirección General de Culturas Populares, D.F., México.
- Torres JC, Fortanelli J, van't Hooft A, Benítez V (2015) **Etnobotánica de la vivienda rural en la región Xi'iuy de la Palma, San Luis Potosí, México.** *Etnobiología* 13:21-36.
- Trotter RT, Needle RH, Goosby E, Bates C, Singer M (2001) **A Methodological Model for Rapid Assessment, Response, and Evaluation: The RARE Program in Public Health.** *Field Methods* 13:137-159.
- Turner N, Deur D, Rae-Mellot C (2011) **“Up on the mointain”: Ethnobotanical importance of montane sites in pacific coastal North America.** *Journal of Ethnobiology* 31(1): 4-43.
- Turvey ST, Pettorelli N (2014) **Spatial congruence in language and species richness but not threat in the world's top linguistic hotspot.** *Proceedings of the Royal Society B* 281: 20141644 doi.org/10.1098/rspb.2014.1644.
- Villaseñor JL (2016) **Checklist of the native vascular plants of Mexico.** *Revista Mexicana de Biodiversidad* 87:559-902.
- Villavicencio MA (2005) **Vegetación e inventario de la flora útil de la Huasteca y la zona Otomí-Tephua de Hidalgo.** SIZA-CONACyT - Universidad Autónoma del Estado de Hidalgo, México.
- Wolverton S, Chambers KJ, Veteto JV (2014) **Climate Change and Ethnobiology.** *Journal of Ethnobiology* 34:273-275.
- Zamudio F, Hilgert N (2012) **Descriptive attributes used in the characterization of stingless bees (Apidae: Meliponini) in rural populations of the Atlantic Forest (Misiones, Argentina).** *Journal Ethnobiology and Ethnomedicine* 8:1-9 doi.org/10.1186/1746-4269-8-9.

Received: 01 February 2019

Accepted: 28 May 2019

Published: 21 June 2019

Supplementary Material

List of biocultural diversity for two communities of the Huasteca, Hidalguense; the type of vegetation is tropical semi-evergreen forest (TSEF) and montane cloud forest (MCF). The table contains the following data: family, gender and species, local name in Spanish and Nahuatl, vegetation type (TSEF / MCF) and category of use and utilization (ceremonial= I, fuel= II, edible= III, IV= construction, V= timber, VI= pet, VII= medicinal, VIII= ornamental, IX= others, X= agricultural use, XI= trade).

Family	Species	Traditional name Spanish/Nahuatl	Type of vegetation	Category of use and utilization
MOLLUSCA				
*Gastropoda	?	caracoles	TSEF/MCF	VII
CRUSTACEA				
*Cambalidae	<i>Procambarus</i> sp.	lunritos / lechuchas	TSEF/MCF	III, X
*Palaemonidae	<i>Macrobrachium</i> sp.	acamayac	TSEF/MCF	III, X
	?	cosoles	TSEF/MCF	III, X
DIPLOPODA				
	?	mipies	MCF	VII
PISCES				
Anguillidae	<i>Anguilla rostrata</i> Lesauur, 1821	coatlmitchi	TSEF	I, III, VII, VIII, X
Lepidosteidae	<i>Lepisosteus osseus</i> Linnaeus, 1758	pez aguja / huitzilmitchi	TSEF	III, VIII, XI
Calosomidae	<i>Ichtiobus labiosus</i> (Meek, 1904)	trampa de pueros / pitzotmitchi	TSEF	III, X
Characidae	<i>Astyanax mexicanus</i> (De Filippi, 1853)	pocta	MCF	III, X
	<i>Cichlasoma labridens</i> (Flegrin, 1900)	mojaras / lecoxtle, xomaquichte, xumaquichte	TSEF	III, VII, X
	<i>Cichlasoma steindachneri</i> (Jordan & Snyder 1898)	mojaras / lecoxtle, xomaquichte, xumaquichte	MCF	III, X
	<i>Hemichthys cyanoguttatus</i> Baird & Girard, 1854	mojaras / lecoxtle, xomaquichte, xumaquichte	MCF	III, X
	<i>Hemichthys tamasopoensis</i> (Amigo Azas, 1993)	mojaras / lecoxtle, xomaquichte, xumaquichte	TSEF/MCF	III, VII, X
Cichlidae	<i>Amatitlania nigrofasciata</i> (Günther, 1857)	mojaras / lecoxtle, xomaquichte, xumaquichte	TSEF/MCF	III, VII, X
	<i>Oreochromis aureus</i> (Steindachner 1864)	tlapia / xumaquichte	TSEF/MCF	III, X
	<i>Oreochromis niloticus</i> (Linnaeus, 1758)	tlapia / xumaquichte	TSEF/MCF	III, X
Cyprinidae	<i>Cyprinella lutrensis</i> G. F. Baird & Girard, 1853	sardina	TSEF	III, X
	<i>Cyprinus carpio communis</i> (Linnaeus, 1758)	carpa / atamichi	TSEF/MCF	III, X
	<i>Dionina ermyzonops</i> Hubbs & Miller 1974	charal / pocta	MCF	III, X
	<i>Dionina igni</i> (Navarez & Navarro 1950)	charal / pocta	TSEF/MCF	III, VII, X
	<i>Dionina rasconis</i> (Jordan & Snyder 1900)	charal / pocta	TSEF	III, VII, X
	<i>Notropis tropicus</i> (Hubbs & Miller, 1975)	charal / pocta	MCF	III, X
Gobiidae	<i>Gobionomus dormitor</i> Lacépède, 1800	huavina, guavina / lecoxtle, tocoxtle	TSEF/MCF	III, X
Ictaluridae	<i>Ictalurus mexicanus</i> (Meek, 1904)	bagra / xoloti, miquacuani, xotomichi	TSEF/MCF	III, VII, X
	<i>Ictalurus punctatus</i> (Constantine Samuel Rafinesque-Schmoltz)	bagra / xoloti, miquacuani, xotomichi	TSEF/MCF	III, X
	<i>Pylaeothus olivaris</i> (Rafinesque, 1818)	bagra / xoloti, miquacuani, xotomichi	TSEF/MCF	III, VII, X
Mugilidae	<i>Agonostomus monticola</i> (Bancroft, 1834)	trucha / itztaomichi	TSEF/MCF	III, VII, X
Poeciliidae	<i>Gambusia vittata</i> Hubbs 1926	charal / pocta	TSEF/MCF	III, X
	<i>Hotonandria bimaculata</i> (Heckel, 1848)	charal / pocta	MCF	III, X
	<i>Poecilia latipunctata</i> (Meek, 1904)	charal / pocta	MCF	III, X
	<i>Poecilia mexicana</i> (Steindachner, 1863)	charal / pocta	MCF	III, X
	<i>Poeciliopsis gracilis</i> (Heckel, 1848)	charal / pocta	MCF	III, X
	<i>Xiphophorus birchmanni</i>	charal / pocta	MCF	III, X
	<i>Xiphophorus confinis</i> (Rauchenberger, Kallman & Menzies, 1990)	charal / pocta	TSEF/MCF	III, X
	<i>Xiphophorus malinche</i> (Rauchenberger, Kallman & Menzies, 1990)	charal / pocta	TSEF/MCF	III, X
	<i>Xiphophorus variatus</i> (Meek, 1904)	charal / pocta	MCF	III, X
AMPHIBIA				
Bufoinae	<i>Incilius nebulifer</i> (Girard, 1843)	ranas ranitas y sapos	TSEF/MCF	VII
	<i>Incilius occidentalis</i> (Camarano, 1879)	ranas ranitas y sapos	TSEF/MCF	VII
	<i>Rhinella marina</i> Linnaeus, 1758	ranas ranitas y sapos	TSEF/MCF	VII
Caudata	?	axoloti / xoloti	TSEF/MCF	
REPTILIA				
Kinosternidae	<i>Kinosternon herrerae</i> Stejneger, 1925	toruga / coxual	TSEF/MCF	II, VI, VII, VIII, XI
Anguillidae	<i>Gerrhonotus ophurus</i> Cope, 1868	escorpión / axitis	TSEF/MCF	
Corytonanidae	<i>Corytophanes hernandezii</i> Wegmann, 1928	laganja verde / iguana	TSEF	VI, XI
Gekkonidae	<i>Hemidactylus frenatus</i> (Schlegel, 1836)	laganja gorda, lujja / axitispitzilziti	TSEF	
Iguanidae	<i>Ctenosaura acanthura</i> (Shaw, 1802)	laganja negra	TSEF	VI, XI
Boidae	<i>Boa constrictor</i> (Linnaeus, 1758)	varacillo / maralcoatl	TSEF/MCF	III, VIII, XI
Colubridae	<i>Coniophanes imperialis</i> (Baird & Girard, 1859)	v-bora de agua	TSEF	

	<i>Drymarchon melanurus</i> (Duméril, Bibron y Duméril, 1854)	vLibora negra / tepeticatl	TSEF/MCF	VIII, XI
	<i>Drymobius margaritiferus</i> (Schlegel, 1837)	chimionera / chacuatla	TSEF/MCF	
	<i>Lampropeltis triangulum</i> (Lacépède, 1788)	coralillo / zicatena	TSEF/MCF	IX
	<i>Spilotes putefus</i> (Linnaeus, 1758)	apachite / apaxcoatl	TSEF/MCF	VII, VIII, XI
	<i>Tantilla rubra</i> (Cope, 1876)	coralillo / zicatena	TSEF/MCF	IX
Dipsosidae	<i>Tropidodipsas sartoni</i> Cope, 1863	coralillo / zicatena	TSEF/MCF	IX
Elapidae	<i>Micurus diastema</i> (Duméril, Bibron y Duméril, 1854)	coralillo / zicatena	TSEF/MCF	IX
Nainiidae	<i>Nerodia rhombifer</i> (Hallowell, 1852)	vLibora de agua	TSEF	
Viperidae	<i>Akistrodon taylori</i> Burger y Robertson, 1951	cola blanca / xochicuicapiili	TSEF/MCF	I, IX
	<i>Atropoides nummifer</i> (Rüppell, 1845)	metlapil / metlapilli	TSEF/MCF	I, IX
	<i>Bothrops asper</i> (Gerron, 1884)	mañuquite	TSEF/MCF	I, VII, IX, XI
	<i>Crotalus</i> sp. Linnaeus, 1758	vLibora de cascabel	TSEF/MCF	I, VII, IX, XI
AVES				
Trogonidae	<i>Crypturellus cinnamomeus</i> (Lesson, 1842)	perdiz / xacoyotl	TSEF/MCF	III, VI, VII, VIII, XI
Anatidae	<i>Carina moschata</i> (Linnaeus, 1758)	pato negro / patox	TSEF/MCF	I, VI, VIII, XI
Crociidae	<i>Ortalis vetula</i> (Wagler, 1830)	chachalaca / acuetlacte	TSEF/MCF	III, VI, VII, VIII, XI
	<i>Penelope purpurascens</i> (Wagler, 1830)	cojilite / cojilote	TSEF/MCF	III, VI, VII, VIII, XI
	<i>Crax rubra</i> Linnaeus, 1758	faisán, paisán, pajarito paisana, pajarito del ocro o pavo real / tepetitotl	TSEF	I, III, VI, VII, VIII, XI
Odonotrogonidae	<i>Dendrotyx barbatus</i> Gould, 1846	godomiz, oodomiz / tzoli	TSEF/MCF	II, VI, VIII, XI
	<i>Colinus virginianus</i> (Linnaeus, 1758)	godomiz, oodomiz / tzoli	MCF	III, VI, VIII, XI
	<i>Dactyortyx thoracicus</i> (Gambel, 1848)	godomiz, oodomiz / tzoli	MCF	III, VI, VIII, XI
Crotonidae	<i>Myctona americana</i> Linnaeus, 1758	ogulula	TSEF	
Pelecanidae	<i>Pelecanus erythrorhynchos</i> Gmelin, 1789	pelicano	TSEF	
Ardeidae	<i>Ardea alba</i> (Linnaeus, 1758)	garza, garza blanca	TSEF/MCF	
	<i>Egretta caerulea</i> (Linnaeus, 1758)	garza negra	TSEF	
	<i>Cochlearius cochlearius</i> (Linnaeus, 1766)	cucharita	TSEF	
Cathartidae	<i>Coragyps atratus</i> (Bechstein, 1783)	zopilote / zapilotl	TSEF/MCF	I, VI
	<i>Cathartes aura</i> (Linnaeus, 1758)	zopilote rojo / zapilotl	TSEF	I, VI
Accipitridae	<i>Fucomis magnirostris</i> (Gmelin, 1768)	colero	TSEF/MCF	VI, VII, VIII, XI
	<i>Buteo platyotus</i> (Schlegel, 1862)	guala, gavilán / cuatcoile	TSEF	VI, VII, VIII, XI
	<i>Buteo jamaicensis</i> (Gmelin, 1788)	guala, gavilán / cuatcoile	TSEF/MCF	VI, VII, VIII, XI
Columbidae	<i>Patagioenas flavirostris</i> (Wagler, 1831)	turcaza, torcaza	TSEF/MCF	III, VI, VII, VIII
	<i>Columba inca</i> (Lesson, 1847)	torolita	TSEF/MCF	VI
	<i>Claravis protiosa</i> (Farran-Pérez, 1886)	paloma	TSEF/MCF	III, VI, VII, VIII
	<i>Leptotilia verreauxi</i> Bonaparte, 1855	paloma	TSEF/MCF	III, VI, VII, VIII
	<i>Zenaidura macroura</i> (Schäfer, 1858)	paloma	TSEF/MCF	III, VI, VII, VIII
	<i>Zenaidura macroura</i> (Linnaeus, 1758)	paloma blanca / singulotl	TSEF/MCF	III, VI, VII, VIII
Tyrtonidae	<i>Tyto alba</i> (Scopoli, 1785)	lechuzca, tecolote lechuzca / tecolotl	TSEF/MCF	I, VII, VIII, XI
Strigidae	<i>Glaucidium sanchezi</i> Lowery & Neuman, 1949	tecolote chico / cuxcux	MCF	I, VII, VIII, XI
	<i>Ciccaba virgata</i> (Cassin, 1849)	tecolote grande / tecolotl	TSEF/MCF	I, VII, VIII, XI
Caprimulgidae	<i>Nyctidromus albicollis</i> (Gmelin, 1768)	tapacaminos / paxuaca	MCF	I, VI
	<i>Antrostomus vociferus</i> (Wils.)	tapacaminos / paxuaca	TSEF	I, VI
Trochilidae	?	chupacucos, colibrí / huitzil	TSEF/MCF	I, VI
Trogonidae	<i>Trogon caligatus</i> Gould, 1835	pajarito amarillo bandera	MCF	VII
	<i>Trogon mexicanus</i> Swainson, 1827	pajarito rojo bandera	MCF	VII
Momotidae	<i>Momotus momota</i> (Linnaeus, 1766)	gorrobano / momot	TSEF/MCF	
Alcedinidae	<i>Megasceryle torquata</i> (Linnaeus, 1766)	marlin pescador	TSEF/MCF	VII
	<i>Chloroceryle amazona</i> (Latham, 1790)	marlin pescador	TSEF/MCF	VII
Ramphastidae	<i>Aulacorhynchus prasinus</i> (Gould, 1834)	tucaneta	TSEF/MCF	I, VIII, XI
	<i>Ramphastos sulfuratus</i> Lesson, 1830	tucán, pito nai, pito nai	TSEF	I, VIII, XI
Picidae	<i>Melanerpes formicivorus</i> (Wagler, 1829)	pajarito carpintero / cuatzurepetl	TSEF/MCF	VII, VIII
	<i>Dryocopus lineatus</i> (Linnaeus, 1766)	pajarito carpintero / cuachenche	TSEF/MCF	VII, VIII
	<i>Campycolaptes curvipennis</i> (Hartlaub, 1844)	pajarito carpintero / cuachenche	TSEF/MCF	VII, VIII
Falconidae	<i>Hierpetotheres cachinnans</i> (Linnaeus, 1758)	guacil	TSEF/MCF	VI, VII, XI
	<i>Micrastur semitorquatus</i> (Vieillot, 1817)	vaquero	TSEF/MCF	VI, VII, XI
	<i>Caracara cheriway</i> (Jacquin, 1784)	huatzi	TSEF/MCF	I, VI, VII, VIII, XI
	<i>Falco sparverius</i> (Linnaeus, 1758)	hacón	TSEF	VI, VII, XI
Psittacidae	<i>Psittacara holochlorus</i> (Sclater, 1858)	colomitos / quilime	TSEF/MCF	III, VI, VII, VIII, XI
	<i>Ficus serilis</i> (Spix, 1824)	perico / cayo	TSEF/MCF	III, VI, VII, VIII, XI
	<i>Amazona viridigenalis</i> (Cassin, 1853)	perico / cutcho	TSEF	III, VI, VII, VIII, XI
	<i>Amazona autumnalis</i> (Lesson, 1842)	perico / cutcho	TSEF	III, VI, VII, VIII, XI
Fumariidae	<i>Xiphorhynchus</i> sp. Swainson, 1827	cuaneztotl	TSEF	
Tyrannidae	<i>Pyrocephalus rubinus</i> (Boddaert, 1783)	cardenal	TSEF/MCF	VI
	<i>Myiozetetes similis</i> (Spix, 1825)	huitlquitzo	TSEF/MCF	
Corvidae	<i>Psittorhynchus morio</i> (Wagler, 1829)	chichiltente	TSEF	
	<i>Cyanocorax yucas</i> (Vigors, 1829)	papán	TSEF/MCF	
	<i>Corvus imparatus</i> Peters, 1929	cuervo / cacalotl	TSEF/MCF	I, VII, XI

Hirundinidae	<i>Hirundo rustica</i> Linnaeus, 1758	golondrina	TSEF/MCF	
Troglodytidae (16);(5)	<i>Myadestes occidentalis</i> Stejneger, 1882 <i>Myadestes unicolor</i> Sclater, 1857 <i>Turdus grayi</i> Bonaparte, 1838	jilguero clarín primavera	TSEF/MCF MCF TSEF/MCF	VI VI VI
Parulidae	?	chikchik / zehuafotometl	TSEF/MCF	
Icteridae	<i>Quiscalus mexicanus</i> (Gmelin, 1786)	tordo / acatzana	TSEF/MCF	I, VII
	<i>Molothrus aeneus</i> (Wagler, 1829) <i>Molothrus ater</i> (Boedaei, 1783) <i>Icterus sp.</i> Bonson, 1780 <i>Psarocolius montezuma</i> (Lesson, 1830) <i>Spinus psaltria</i> (Say, 1823)	tordo de ojos rojos / pipix tordo / acatzana calandria, pajarito del sol / tonaitototl papón real dominico	TSEF/MCF TSEF/MCF TSEF/MCF TSEF/MCF TSEF/MCF	I, VII I, VII VII VII
MAMMALIA				
Didelphidae	<i>Didelphis marsupialis</i> Linnaeus, 1757 <i>Didelphis virginiana</i> Kerr, 1792 <i>Philander opossum</i> (Linnaeus, 1758) <i>Marmosa mexicana</i> Merriam, 1897	tlacuache / tlacuahuilotl tlacuache grade / itztatlacuahuilotl zhuantale raí'in tlacuache / quimichitlacuahuilotl	TSEF/MCF TSEF/MCF TSEF/MCF TSEF/MCF	III, VI, VII, VIII III, VI, VII, VIII VII
Dasyproctidae	<i>Dasyprocta novemcinctus</i> Linnaeus, 1758	armadillo / atitche	TSEF/MCF	I, III, VII
Myrmecophagidae	<i>Tamandua mexicana</i> (de Sausure, 1860)	oso mielero, oso horniguero / xopa miel	TSEF/MCF	VII, VIII
Orden Soricomorpha	?	raí'in / quimichi	TSEF/MCF	
Orden Chiroptera	?	murciélago / calzoath, tzotzo	TSEF/MCF	I, VII
**Atelinae	** <i>Atelopus geoffroyi</i> Kuhl, 1820	*mono	MCF	
Leporidae	<i>Sylvilagus brasiliensis</i> (Linnaeus, 1758)	conejo / cuatochi	TSEF/MCF	III, VI, VII, VIII, IX, XI
Sciuridae	<i>Sciurus aureogaster</i> (Cuvier, 1829) <i>Sciurus deppii</i> (Perles, 1864)	ardilla, ardilla colorada / tecomate ardilla, ardilla negra / tecomate	MCF TSEF/MCF	III, VI, VII, VIII III, VI, VII, VIII
Geomysidae	<i>Orthogeomys hispidus</i> (Le Conte, 1852)	tuza, tuza de tierra, tuza raicera / tlaltuza	TSEF/MCF	III, IX
Erethizontidae	<i>Coscodon mexicanus</i> (Kerr, 1792)	puercito espín / xompi	TSEF/MCF	III, VI, VII, VIII, XI
Cuniculidae	<i>Cuniculus paca</i> (Linnaeus, 1776)	tuza real / cuatuzo, tuza real	TSEF/MCF	III, VI, VII, VIII, XI
Cnecidae	?	raí'in / quimichi	TSEF/MCF	
Felidae	<i>Herpailurus yagouaroundi</i> (Leclercq, 1809) <i>Leopardus pardalis</i> (Linnaeus, 1758) <i>Leopardus wiedii</i> (Schinz, 1821) <i>Lynx rufus</i> (Schreber, 1777) <i>Puma concolor</i> (Linnaeus, 1771) <i>Panthera onca</i> (Linnaeus, 1758)	onza / sacamixtle ocelote, leoncillo / ocelotl tigriño / cuamotamixto gato montés, gato de monte, gatillo / cuamotamixto llín / tecuari tigre / tecuaquetl, tecuari	TSEF/MCF TSEF/MCF TSEF/MCF TSEF/MCF TSEF/MCF TSEF/MCF	VII, VIII, XI VI, VII, VIII, XI III, VI, VII, VIII, XI VI, VII, VIII, XI VIII, XI VIII, XI
Canidae	<i>Canis latrans</i> Say, 1823	coyote, perro coyote / coyotl, coyachichi	TSEF/MCF	I, III, VII, VIII, IX, XI
	<i>Urocyon cinereoargenteus</i> (Schreber, 1775)	zorro / coyochi, itzoayuchi	TSEF/MCF	III, VII, VIII, IX, XI
Mephitidae	<i>Conepatus leucocotus</i> (Lichtenstein, 1832) <i>Mephitis macroura</i> Lichtenstein, 1832	zorillo / epa zorillo / epa	TSEF/MCF TSEF/MCF	III, VII, VIII, XI III, VII, VIII, XI
Mustelidae	<i>Spilogale gracilis</i> Merriam, 1890 ** <i>Lontra longicaudis</i> (Olfers, 1818) <i>Eira barbara</i> (Linnaeus, 1758) <i>Galliois vittata</i> (Schreber, 1776) <i>Mustela frenata</i> Lichtenstein, 1831	zorillo / epa *perro de agua / atlochich perro de cenzo, perro de monte / tepachichi grisón hurón	TSEF/MCF MCF TSEF/MCF TSEF/MCF	III, VII, VIII, XI I, VIII, XI VII VI, VIII
Procyonidae	<i>Potos flavus</i> (Schreber, 1774) <i>Nasua narica</i> (Linnaeus, 1776) <i>Procyon lotor</i> (Linnaeus, 1758)	mapache / tancho rejón / pezocle mapache / coaticuamixto, mapachi	TSEF/MCF TSEF/MCF TSEF/MCF	III, VI, XI III, VI, VII, VIII, XI III, VI, VII
Tayassuidae	<i>Dicotyles angulatus</i> (Cope, 1869)	jabalí / coapizotl	TSEF/MCF	III, VI, VIII, XI
Cervidae	<i>Mazama temama</i> (Kerr, 1792) <i>Odocoileus virginianus</i> (Zimmermann, 1780)	venado chico / cuachacal, chacal venado / mazatl	TSEF/MCF TSEF/MCF	III, VI, VIII, XI III, VI, VII, VIII, XI
FUNGI				
Auriculariaceae	<i>Auricularia auricula</i> (L.: Fr.) Lindau <i>Auricularia delicata</i> (Fr.) Henn. <i>Auricularia polytricha</i> (Monn.) Sacc.	oreja, oreja de viejita, nalga de la abuelita / cuapofachiquinte, cuafolashitl oreja, oreja de viejita, nalga de la abuelita / cuapofachiquinte, cuafolashitl oreja, oreja de viejita, nalga de la abuelita / cuapofachiquinte, cuafolashitl	TSEF/MCF TSEF/MCF TSEF/MCF	III III III
Lycoperdaceae	<i>Calvatia excipuliformis</i> (Pers: Pers.) Penderg	peporitas, huevos de venado / mazatltequistle, mazatlmenacal	MCF	III
Pleurotiaceae	<i>Pleurotus djamour</i> (Fr.) Boedj. <i>P. albidus</i> (Berk.) Pegler	hongro blanco / totomoshchiquinte, totomoshtrianacal hongro palón blanco / itzlecnenacal	TSEF/MCF TSEF/MCF	III, XII III, XII
Schizophyllaceae	<i>Schizophyllum commune</i> Fr.	chiquinte	TSEF/MCF	

Cantharellaceae	<i>Cantharellus cibarius</i> Fr.	hongos amarillos - amarillo chiquinte / <i>xochilnanacatl, chipahuachiquinte,</i>	TSEF/MCF	III, XII
	<i>Cantharellus friesii</i> Peck	hongos amarillos - amarillo chiquinte / <i>xochilnanacatl, chipahuachiquinte</i>	TSEF/MCF	III, XII
	<i>Cantharellus latentius</i> (Berck.) Singer	hongos amarillos - amarillo chiquinte / <i>xochilnanacatl, chipahuachiquinte</i>	TSEF/MCF	III, XII
	<i>Cantharellus lewissii</i>	moradito, moradito chiquinte / <i>xochilnanacatl, xocoyochiquinte</i>	TSEF/MCF	III, XII
	<i>Cantharellus minor</i> Peck	hongos amarillos - amarillo chiquinte / <i>xochilnanacatl, chipahuachiquinte</i>	TSEF/MCF	III, XII
Polyporaceae	<i>Lentinus</i> sp. Fr.	flanchino/chiquinte	TSEF	III
Ustilaginaceae	<i>Ustilago maydis</i> (D.C.) Corda	hongo de milpa, hongo de maíz / <i>cuibaccoche</i>	TSEF/MCF	III
PLATAE				
Acanthaceae	<i>Justicia spicigera</i> Schldl.	mahuite, mohuate	TSEF/MCF	VII
Actinidiaceae	<i>Saurauia scabrifolia</i> Hemsl.	acalama	TSEF/MCF	
Agavaceae	<i>Agave</i> sp. L. 1753	Maguey	TSEF/MCF	
	<i>Yucca</i> sp. L.	isole cimarrón	TSEF/MCF	VII
Amaranthaceae	<i>Amaranthus hybridus</i> L.	quelita / <i>quilitl</i>	TSEF/MCF	III
	<i>Amaranthus acanthochiton</i> J.D. Sauer	ameranto	TSEF	
	<i>Celosia cristata</i> L.	mano de león	TSEF/MCF	I, VIII, XI
	<i>Gomphrena globosa</i> L.	diante de león	TSEF/MCF	I, VIII, XI
Amaryllidaceae	<i>Allium glandulosum</i> Link & Otto.	cebollita / <i>xonacato, xonacatl</i>	TSEF/MCF	III, VII, XI
Anacardiaceae	<i>Mangifera indica</i> L.	mango criollo, mango manila, mango petacón	TSEF/MCF	II, III, VIII, X, XI
	<i>Spondias mombin</i> L.	jobo / <i>cuaxocotl</i>	TSEF/MCF	II, III, VII, X, XI
	<i>Spondias purpurea</i> L.	chuela	TSEF/MCF	II, III, VII, X, XI
Annonaceae	<i>Annona cherimola</i> Mill.	anona	TSEF/MCF	II, III, V, XI
	<i>Annona muricata</i> L.	guanabana	TSEF	II, III, V, XI
	<i>Rollinia membranacea</i> Triana & Planch.	anona	TSEF	II, III, V, XI
Apiaceae	<i>Conandrum</i> sp. L.	clantró / <i>colantró</i>	TSEF/MCF	III, XI
	<i>Eryngium foetidum</i> L.	clantró extranjero / <i>viscotantró</i>	TSEF/MCF	III, XI
	<i>Foeniculum vulgare</i> Mill.	finjo	TSEF/MCF	III, VII, XI
Apocynaceae	<i>Allamanda</i> sp. L.	copa de oro, copa de rey	TSEF/MCF	VII
	<i>Plumeria acutifolia</i> Poirai	flor de mayo, flor de la cruz / <i>cacalo/xuchitl</i>	TSEF/MCF	I, II, VII, VIII, XI
	<i>Plumeria rubra</i> L.	flor de mayo, flor de la cruz / <i>cacalo/xuchitl</i>	TSEF/MCF	I, II, VII, VIII, XI
	<i>Stemmadenia donnell-smithii</i> (Rose) Woods	cojón de gato	TSEF/MCF	VII, XI
	<i>Thevetia peruviana</i> A. D.C.	huevos de toro	TSEF	VII, VIII
Araceae	<i>Syngronium podophyllum</i> Schott.	chapas	TSEF/MCF	VII, VIII
	<i>Xanthosoma robustum</i> Schott.	hoja de luna / <i>quoquesquilitl</i>	TSEF/MCF	III, VIII
	<i>Zantedeschia aethiopica</i> (L. Spreng.)	alcarráz	MCF	I, VIII, XI
Araliaceae	<i>Dendropanax arboreum</i> (L.) Don. & Planch.	tamalcóhuitl	TSEF/MCF	II, IV, V, XI
Arecaceae	<i>Acrocomia aculeata</i> (Jacq.) Lodd. Ex Mart	coyol	TSEF	III, XI
	<i>Chamaedorea elegans</i> Mart	palmita	TSEF	I, VIII, XI
	<i>Chamaedorea liebmanni</i> Martius	palmita	MCF	I, VIII, XI
Asclepiadaceae	<i>Asclepias curassavica</i> L.	venanillo, quebra huesos	TSEF/MCF	VII
	<i>Gonolobus niger</i> R.Br.	apayote / <i>cuahuayote</i>	TSEF/MCF	
Aspleniaceae	<i>Asplenium sphaerosporum</i> A.R. Sm.	pesma chico, pesmita	MCF	
Asieraceae	<i>Achillea millefolium</i> L.	plumajillo	MCF	VII, VIII
	<i>Ageratum corymbosum</i> Zucc. Ex Pers.	hierba del pasino	MCF	VII, VIII
	<i>Aldama dentata</i> La Llave & Lex.	acahual	TSEF/MCF	IX
	<i>Artemisa</i> sp. L. 1753	quenjo	TSEF/MCF	VII
	<i>Artemisia ludoviciana</i> Nutt.	estafate	MCF	VII
	<i>Baccharis conferta</i> Kunth	escobilla / <i>tepetlapushitl</i>	TSEF/MCF	IX
	<i>Bidens pilosa</i> L.	mozotl	TSEF/MCF	IX
	<i>Matricaria recutita</i> L.	manzanilla	TSEF/MCF	VII, XI
	<i>Senecio confusus</i> Britton	pinca	TSEF/MCF	VII
	<i>Porophyllum ruderale</i> (DC.) Cronquist	papalo quelele	TSEF/MCF	III, XI
	<i>Tagetes coronopifolia</i> Willd.	oemopal silvestre / <i>compoal/xuchitl</i>	TSEF/MCF	I
	<i>Tagetes erecta</i> L.	oemopal / <i>compoal/xuchitl</i>	TSEF/MCF	I, VII, VIII, XI
Begoniaceae	<i>Begonia heracleifolia</i> Schldl. & Cham.	concella / <i>mayahual</i>	MCF	
	<i>Begonia nelumbifolia</i> Schldl. & Cham	bandejita / <i>mayahual del maco</i>	MCF	
	<i>Begonia wallichiana</i> Lahm	xihuita amargoso	MCF	
Bignoniaceae	<i>Crescentia cujete</i> L.	timo / <i>cuatecomate, cuatecomactl</i>	TSEF/MCF	IX
	<i>Jacaranda mimosaeifolia</i> D. Don	jacaranda	TSEF/MCF	VII
	<i>Panmentiera aculeata</i> (Kunth) Seemann	chole	TSEF/MCF	II, III, VII
	<i>Spathodea campanulata</i> Beauv.	San Joséxuchitl	TSEF/MCF	I, VIII
	<i>Tabebuia rosea</i> (Berthel.) A. D.C.	palo de rosa	TSEF	II, VIII
	<i>Tocoma stans</i> (L.) HBK.	<i>ahuaxuchibara</i>	TSEF/MCF	
Bombacaceae	<i>Coiba pontandra</i> (L.) Gaert.	osiba	TSEF/MCF	II, IV, V, XI
	<i>Pseudobombax ellipticum</i> (Kunth) Dugand.	cabello de ángel / <i>tzotzocole</i>	TSEF	II, IV, VIII

Brassicaceae	<i>Brassica</i> aff. <i>Oleracea</i> L.	quelite de col / <i>colquiliti</i>	TSEF/MCF	III	
Bromeliaceae	<i>Aechmea bracteata</i> (Sw.) Griseb.	amoxco, tecolomefi	TSEF		
	<i>Ananas comosus</i> L. Merr.	pica	TSEF/MCF	III	
	<i>Tillandsia recurvata</i> (L.) L. 1762	timbinche	TSEF/MCF		
Burseraceae	<i>Tillandsia usneoides</i> (L.) L.	ieno	MCF	I, VIII	
	<i>Bursera paniculata</i> (Sessl & Moc.) ex DC. Engel	palo brujo	TSEF	II, IV	
	<i>Bursera simaruba</i> (L.) Sarg	chaca	TSEF/MCF	II, IV, VII	
Cactaceae	<i>Protium copal</i> (Schltdl. & Cham.) Engl.	copal	TSEF/MCF	I, II, IV, X, XI	
	<i>Acanthocereus</i> sp. (Engelm ex A. Berger) Britton & Rose	rana-nopal / <i>jacube</i>	TSEF/MCF	III, XI	
	<i>Hyloteleium undatum</i> (Haw.) Britton & Rose	piaya	TSEF/MCF	III, XI	
Caprifoliaceae	<i>Opuntia streptacantha</i> Lam.	nopal	TSEF/MCF	II, XI	
	?	ligano	TSEF	VII	
Cannaceae	<i>Sambucus mexicana</i> Presl.	saúco	MCF	II, X	
Caryophyllaceae	<i>Canna cauliflora</i> Jacq.	ochonito, ochonictli	TSEF/MCF	III	
	<i>Canna papaya</i> L.	papaya	TSEF/MCF	II, XI	
Cecropiaceae	<i>Cecropia obtusifolia</i> Bertol.	palo de canal, palo chiflón, hormiguillo / <i>cuauquiquistli</i>	TSEF/MCF	II, IV, VII	
Chenopodiaceae	<i>Chenopodium ambrosioides</i> L.	epazote / <i>epazotli</i>	TSEF/MCF	III, VII, XI	
Clethraceae	<i>Clethra mexicana</i> DC.	aguacatillo	MCF		
Combratiaceae	<i>Clethra pringlei</i> S. Watson	aguacatillo	MCF		
	<i>Terminalia catappa</i> L.	almendro	TSEF/MCF	II, VIII, X	
Convolvulaceae	<i>Ipomoea batatas</i> Lam.	camote	TSEF/MCF	III, XI	
	<i>Ipomoea purga</i> (Wender.) Hayne	zoys	TSEF/MCF	III	
Crasulaceae	<i>Ipomoea</i> sp. Senth. Ex. Vell.	bejuco de flores azules, quiebra platos	TSEF/MCF		
	<i>Kalanchoe pinnata</i> (Lamark) Pearson	tronadora	TSEF/MCF	IX	
Cucurbitaceae	<i>Citrullus lanatus</i> (Thunb.) Matsum. & Nakai 1916	sandía	TSEF	III, XI	
	<i>Cucurbita maxima</i> Lam.	calabaza	TSEF/MCF	III, XI	
	<i>Seschium edule</i> (Jacq.) Siu.	chayote	TSEF/MCF	III, XI	
Chrysobalanaceae	<i>Couepia polyandra</i> (Kunth) Rosa	olopillo	TSEF		
Cyatheaceae	<i>Cyathea fulva</i> (M. Martens & Galeotti)	pecina grande	MCF	VII	
Cyperaceae	<i>Cyperus odoratus</i> L.	zacate	TSEF/MCF	IX	
Ebenaceae	<i>Diospyros digyna</i> Jacq. Pl. hort. schoenbr.	zapota negro / <i>zapoti</i>	TSEF/MCF	III, X, XI	
Euphorbiaceae	<i>Cnidioscolus acornifolius</i> (Mill.) I. M. Johnston.	chaya / <i>chichacuahuatl</i>	TSEF/MCF		
	<i>Cnidioscolus multiobus</i> (Pax) I.M. Johnston	tesonquiliti	TSEF/MCF		
	<i>Croton draco</i> Schltdl.	sangracao / <i>escuahuitl</i>	TSEF/MCF	I, VII	
	<i>Croton pulcher</i> Müll. Arg.	soliman	TSEF/MCF		
	<i>Euphorbia pulcherrima</i> Willd. ex Klotzsch	flor de la virgen, noche buena / <i>tonantzínkuhuatl</i>	TSEF/MCF	I, VIII, XI	
	<i>Jatropha curcas</i> L. 1753	pico	TSEF/MCF	II	
	<i>Marihot esculenta</i> Crantz	yuca	TSEF/MCF	III, XI	
	<i>Ricinus communis</i> L.	iguera	TSEF/MCF		
	Equiceae	<i>Equisetum</i> sp. L.	cola de caballo / <i>aquiztopile</i>	TSEF/MCF	III
	Fabaceae	<i>Acacia corrifera</i> L. Willd.	toro cuamo	TSEF/MCF	II
		<i>Acosmium panamense</i> (Benth.) Yakovlev	hueso de tigre, huasillo	TSEF	II, IV
		<i>Amicia zygomera</i> DC.	papaloxhuatl	TSEF	
		<i>Arachis hypogaea</i>	cacahuata	TSEF/MCF	III
		<i>Bauhinia divaricata</i> L.	pata de buey	TSEF	I, IV, VIII
		<i>Caesalpinia pulcherrima</i> Benth.	unco de rosal / <i>tabachín</i>	TSEF/MCF	VII
<i>Cajanus cajan</i> (L.) Willsp.		lentejilla, frijol grande	TSEF/MCF	III	
<i>Cassia fistula</i> L.		lluvia de oro	TSEF/MCF	II, IV, VIII	
<i>Calliandra grandiflora</i> (L.) Hitchc. Benth.		cabello de ángel / <i>tzotzocole</i>	TSEF/MCF		
<i>Dalbergia palo-escrito</i> Rzed. & Gündl-Gómez		palo escrito	TSEF/MCF	IV, V, XI	
<i>Delonix regia</i> (Bojer) Raf.		framboyán / flor de mayo / flor de las madres	TSEF/MCF	II, IV, VIII	
<i>Diphysa senoides</i> Benth.		quebra hada / <i>quebrache, quebrachi</i>	TSEF/MCF	IV, V	
<i>Erythrina americana</i> Mill.		penuche / <i>penuch</i>	TSEF/MCF	II, III, IV, VII, IX	
<i>Erythrina lanata</i> Rose		penuche / <i>penuch</i>	TSEF/MCF	II, IV, VII, IX	
<i>Eysenhardtia platycarpa</i> Pennell & Gaff.		palo azul	TSEF	VII	
<i>Clinidialia sepium</i> (Jacq.) Steud.	cacahuatl	TSEF			
<i>Mucuna pruriens</i> (L.) DC.	pica pica	TSEF/MCF			
<i>Phaseolus coccineus</i> L.	frijol	TSEF/MCF	III, XI		
<i>Phaseolus</i> sp. L.	cashitán	TSEF/MCF	III		
<i>Pisoidia piscipula</i> (L.) Sarg.	chijol	TSEF			
<i>Pithecolobium dulce</i> (Roxb.) Benth.	frijolillo / humo	TSEF/MCF	III, XI		
<i>Tamarindus indica</i> L.	temerindo / <i>tamarindocuahuitl</i>	TSEF/MCF	II, III, IV, XI		
<i>Trifolium</i> sp. L.	quelite estrella	TSEF/MCF			
<i>Vicia faba</i> L.	haca	TSEF/MCF	III, XI		
Fagaceae	<i>Quercus affinis</i> Scheidtl.	encino / <i>ticcuahuatl</i>	MCF	II, IV, V, XI	
	<i>Quercus coccinea</i> (L.) Liebm.	encino / <i>ticcuahuatl</i>	MCF	II, IV, V, XI	
	<i>Quercus oleoides</i> Cham & Schltdl.	encino / <i>ticcuahuatl</i>	TSEF	II, IV, V, XI	
	<i>Quercus sartorii</i> N.Ze	encino / <i>ticcuahuatl</i>	MCF	II, IV, V, XI	
	<i>Quercus xalapensis</i> Humb. & Bonpl.	encino / <i>ticcuahuatl</i>	MCF	II, IV, V, XI	

Gramineae	<i>Lasiacis divaricata</i> (L.) A. S. Hitch.	bambú	TSEF	II, IV
Hamamelidaceae	<i>Liquidambar macrophylla</i> Oerst.	suchiata / suchiatl	TSEF/MCF	II, IV
Heliconiaceae	<i>Heliconia collinsiana</i> Griggs.	papatla	MCF	III, VIII
	<i>Heliconia schiedeana</i> Klotzsch.	papatla	TSEF	III, VIII
Lamiaceae	<i>Agastache mexicana</i> (Kunth) Linton et Epling	ironjil	MCF	VII
	<i>Clerodendrum thomsoniae</i> Balf.	velo de novia	TSEF	VIII, IX
	<i>Meritha sprucei</i> L.	hierba buena	TSEF/MCF	VII
	<i>Ocimum basilicum</i> L.	albahaca	TSEF/MCF	I, III, VII
	<i>Origanum majorana</i> L.	mejorana	TSEF/MCF	VII
	<i>Origanum vulgare</i> L.	orégano	TSEF/MCF	III, VII, XI
	<i>Rosmarinus officinalis</i> L.	romero	TSEF/MCF	I, VII
	<i>Salvia leucantha</i> Cav.	cordoncillo	MCF	
	<i>Salvia mexicana</i> L.	flanchichinole	MCF	
	<i>Sesamum indicum</i> L.	ajonjolí	TSEF/MCF	III, XI
	<i>Stachys parvifolia</i> Martens & Galeotti	talachia	MCF	
Lauraceae	<i>Cinnamomum camphora</i> L.	alcanfor	TSEF/MCF	VII
	<i>Cinnamomum verum</i> J. Presl	canela	TSEF/MCF	III
	<i>Licania capitata</i> (Cham. & Schltdl.) Kosterm.	xotime	MCF	
	<i>Persea americana</i> Miller	aguacate / aguacatl	TSEF/MCF	II, III, IV, VII, X, XI
	<i>Persea americana</i> var. <i>drymifolia</i> (Schltdl. & Cham.) S. F. Blake	aguacate oloroso / aguacatl tlapanilí	TSEF/MCF	II, III, IV, VII, X, XI
	<i>Persea schiedeana</i> Nees	patula / pahuatl	TSEF/MCF	II, III, IV, VII, X, XI
Liliaceae	<i>Allium sativum</i> L.	ajo	TSEF/MCF	III, VII
Lophosoriaceae	<i>Lophosoria quadriripinata</i> (J. F. Gmel.) C. Chr.	pesma	MCF	
Loranthaceae	<i>Struthanthus crassipes</i> (Oliver) Eichler	seca palo	TSEF/MCF	
Magnolaceae	<i>Magnolia grandifolia</i> L.	magnolia	TSEF	VII
	<i>Talauma mexicana</i> (DC.) Don	yoloxuchitl	TSEF	VI, VIII
Malpighiaceae	<i>Burchardia palmieri</i> S. Watson	huasha	TSEF/MCF	
	<i>Byrsonima crassifolia</i> (L.) Kunth	nantzi	TSEF	II, III, XI
Malvaceae	<i>Hibiscus rosa-sinensis</i> L.	tulipán	TSEF/MCF	VII
	<i>Hibiscus sabdariffa</i> L.	jamaica	TSEF/MCF	III, VII, XI
	<i>Malvastrum arboreum</i> Cav.	manzanilla	TSEF/MCF	VII
	<i>Quararíbea funebris</i> (La Llave) Vischer	cacahuaxochitl	TSEF/MCF	
Melastomataceae	<i>Conostegia xalapensis</i> (Bonpl.) D. Don ex DC.	festiva	MCF	
	<i>Arthrostemma ciliatum</i> Pav. ex D. Don	quelite agrio / xocoquilitl	TSEF/MCF	III
Meliaceae	<i>Azadirachta indica</i> A. Juss.	nin	TSEF	
	<i>Cedrela odorata</i> L.	cedro, cedro rojo / ticuahuítl	TSEF/MCF	IV, V, XI
	<i>Guarea glabra</i> Vahl	palo blanco / itztacuahuitl	TSEF	
	<i>Melia azedarach</i> L.	proche, prodri	TSEF/MCF	II, IV
	<i>Swietenia macrophylla</i> King	caoba	TSEF/MCF	IV, V, XI
Menispermataceae	<i>Ossampelos owariensis</i> Beauvais ex DC.	arifzo	MCF	
Mimosaceae	<i>Calliandra grandiflora</i> (L.) H.Dr. Benitt.	cabello de Engel / tzotzoote	TSEF/MCF	VII
Mimosaceae	<i>Inga jinicuil</i> Schltdl. & Cham. Ex G. Don	chalahuite / chalahuitl	TSEF/MCF	II, IV
Moraceae	<i>Inga vera</i> Willd.	chalahuite / chalahuitl	TSEF/MCF	II, IV
	<i>Brosimum alicastrum</i> Sw.	qijte	TSEF/MCF	
	<i>Castilla elastica</i> Cav.	palo de hule / olicuahuítl	TSEF	IV
	<i>Ficus padifolia</i> H.B.K.	jalamaire / xalamati	TSEF/MCF	
	<i>Macclura tinctoria</i> (L.) D. Don ex Steud	mora	TSEF/MCF	II, IV, V
Muntingiaceae	<i>Muntingia calabura</i> L.	capullín / pujhua	TSEF	III
Musaceae	<i>Musa paradisiaca</i> L.	plátano	TSEF/MCF	III, XI
	<i>Musa balbisiana</i> Colla	plátano macho	TSEF/MCF	III, XI
Myrsinaceae	<i>Myrica centifera</i> L.	ahuaxochitl	MCF	
	<i>Eucalyptus</i> sp. L. Her.	eucalipto	MCF	
Myrsinaceae	<i>Stylogyne</i> sp. A. DC.	capullín / capollín	TSEF	III
Myrtaceae	<i>Eugenia capuli</i> (Cham. & Schltdl.) O. Berg	pieste	TSEF	
	<i>Pimenta dioica</i> (L.) Merr.	pimienta	TSEF/MCF	II, III, IV, V, X, XI
	<i>Psidium guajava</i> L.	guayaba / chaltchocotl	TSEF/MCF	II, III, VII, X, XI
Myrtaceae	<i>Bougainvillea glabra</i> Choisy	bugambilia	TSEF/MCF	VII
Onagraceae	<i>Fuchsia microphylla</i> Kunth	aretillo	TSEF/MCF	
Orchidaceae	<i>Oncidium sphaelatum</i> Lindl.	flor de la santa cruz, flor de mayo, flor de cuaresma	TSEF/MCF	I, VIII
	<i>Vanilla planifolia</i> Jacke.	vanilla	TSEF	III, VIII, XI
Oxalidaceae	<i>Averrhoa carambola</i> L.	carambola	TSEF	III, XI
Pinaceae	<i>Pinus greggii</i> Englem.	pino	MCF	II, IV
	<i>Pinus patula</i> Schlecht et Cham.	pino	MCF	II, IV
Piperaceae	<i>Piper auritum</i> Kunth	tequellie / acoyo, tequiquilitl	TSEF/MCF	III
	<i>Peperomia</i> sp. Ruiz & Pav.	verdolaga de monte	TSEF/MCF	III
Platanaceae	<i>Platanus mexicana</i> Moric.	lilamo / acuahuitl	TSEF	II, IX
Poaceae	<i>Arundo donax</i> L.	carizo	TSEF/MCF	IV, IX
	<i>Echinochloa colona</i> (L.) Link	zacate colorado	TSEF/MCF	IX
	<i>Digitaria decumbens</i> Steud.	pasto pangola	TSEF/MCF	IX
	<i>Guadua amplexifolia</i> J. Presl	olafe	TSEF/MCF	II, IV, IX
	<i>Pennisetum purpureum</i> Schumacher.	pasto talluín	TSEF/MCF	IX
	<i>Saccharum officinarum</i> L.	caña de azúcar	TSEF/MCF	II, III, IX, XI

Polypodiaceae	<i>Phlebodium aureum</i> (L.) J. Smith	ostilla de león / <i>tehuancachitlauhuitl</i>	MCF	
	<i>Rumex</i> sp. L.	lengua de vaca	TSEF/MCF	III
Pteridaceae	<i>Adiantum</i> sp. L.	pesmita	TSEF/MCF	
Rhamnaceae	<i>Karwinskia humboldtiana</i> (Roemer & Schultes) Zucc.	capulín de zorra, capulín amarillo	TSEF	III
Rosaceae	<i>Eriobotrya japonica</i> (Thunb.) Lindl.	anisero, niñero	MCF	III
	<i>Prunus persica</i> (L.) Batsch	durazno	TSEF/MCF	III
	<i>Prunus serotina</i> Ehrh.	capulín	TSEF/MCF	III
Rubiaceae	<i>Coffea arabica</i> L.	café	TSEF/MCF	III, VII, IX, XI
	<i>Hamelia patens</i> Jacq.	maduro zapote / <i>cacahuaxochitl</i>	TSEF/MCF	VII
	<i>Galium mexicanum</i> Kunth	pegaropa	MCF	
Rutaceae	<i>Casimiroa edulis</i> Llave & Lex.	zapote blanco / <i>iztaczapote</i>	TSEF	
	<i>Citrus aurantiifolia</i> (Christm.) Swingle	naranja cimanona	TSEF/MCF	III
	<i>Citrus sinensis</i> Osbeck	naranja	TSEF/MCF	III, IX, XI
	<i>Citrus limetta</i> Risso	lima	TSEF/MCF	III
	<i>Citrus maxima</i> (Burm. ex Rumph.) Merr.	limón	TSEF/MCF	III, VII, XI
	<i>Citrus reticulata</i> Blanco	mandarina	TSEF/MCF	III, IX, XI
	<i>Citrus x paradisi</i> Macfad.	toronja	TSEF/MCF	III
	<i>Munaya paniculata</i> Jokoening Ex L.	limonana	TSEF	VII
	<i>Ruta chalepensis</i> L.	ruda	TSEF/MCF	VII
Salicaceae	<i>Salix humboldtiana</i> Willd.	saúce	TSEF	X
	<i>Zuelania guidonia</i> (Sw.) Britt. & Millsp.	volarín	TSEF	I
Sapindaceae	<i>Litchi chinensis</i> Sonn.	lichi	TSEF/MCF	III, XI
	<i>Sapindus saponaria</i> L.	jaborcito	TSEF	IX
Sapotaceae	<i>Marilkera zapota</i> (L.) van Royen	chicozapote / <i>xicozapote</i>	TSEF/MCF	II, III, X, XI
	<i>Pouteria campechiana</i> (Kunth) Baehni	zapote amarillo	TSEF/MCF	III
	<i>Pouteria sapota</i> (Jacq.) H.E. Moore & Stearn.	zapote mamey / <i>zapote</i>	TSEF/MCF	II, III, IV, VII, X, XI
Simaroubaceae	<i>Simarouba glauca</i> DC.	piñache	TSEF	II, IV
Solanaceae	<i>Capsicum annuum</i> L.	chilepín	TSEF/MCF	III
	<i>Capsicum</i> sp. L.	chile	TSEF/MCF	III
	<i>Brugmansia</i> sp. <i>candida</i> Pers.	<i>xochilcampana</i>	TSEF/MCF	VII, VIII
	<i>Nectandra sanguinea</i> Roitb.	aguacachile	TSEF/MCF	
	<i>Nicotiana tabacum</i> L.	tabaco	TSEF/MCF	I, VII, IX, XI
	<i>Solanum nigrescens</i> Mart. & Gal.	tomatillo	TSEF/MCF	III, XI
Sterculiaceae	<i>Guazuma ulmifolia</i> Lam.	guasima / <i>aguiche</i>	TSEF/MCF	II
	<i>Theobroma cacao</i> L.	cacao	TSEF	III, VIII
Tiliaceae	<i>Helicarpus donnell-smithii</i> Rose	jonota	TSEF/MCF	X
Ullinaceae	<i>Aphananthe monoica</i> (Heindl.) J.F. Leroy	ajuate, <i>ajuactli</i>	TSEF	
	<i>Ulmus mexicana</i> (Liebm.) Planch.	palo varín, palo de hombre / <i>tlacacuahuitl</i>	TSEF/MCF	IV, V
	<i>Troma micrantha</i> (L.) Blume	gunda, capulín	TSEF	III
Urticaceae	<i>Urtica caracasana</i> (Jacq.) Griseb.	ortiga real	TSEF/MCF	VII
	<i>Urtica mexicana</i> Liebm.	ortiga	TSEF/MCF	VII
Verbenaceae	<i>Petrea volubilis</i> L.	raspasombreiro	TSEF/MCF	II, III, VII, VIII
Vitaceae	<i>Vitis filifolia</i> Humb. & Bonpl. ex Roemer & Schult.	mecate de uva / <i>cuaxocomecati</i>	TSEF/MCF	III
Zamiaceae	<i>Zamia heimerlæ</i> Calderín & Standl.	zompollo	TSEF/MCF	
Zingiberaceae	<i>Costus mexicanus</i> Liebm.	cañal de jabalí / <i>cuapitzotl</i> , <i>acaticuapitzihuitl</i>	TSEF/MCF	VII
	<i>Zingiber officinale</i> (L.) Roscoe.	jengibre	TSEF/MCF	III, VII, XI