



Conflicts between humans and crocodilians in urban areas across Brazil: a new approach to support management and conservation

Paulo Braga Mascarenhas-Junior^{1,9*}, Fábio Maffei^{2,9}, Fábio Muniz^{3,9},
Ricardo F. Freitas-Filho^{4,5}, Thiago Costa Gonçalves Portelinha^{6,9},
Zilca Campos^{7,9} and Luís A. B. Bassetti^{8,9}

ABSTRACT

Crocodilian-human conflicts, caused especially by urban expansion and habitat destruction, have been considered one of the main threats to the conservation of these species worldwide. In Brazil, such conflicts have been documented with crocodilian species all over the country. This study identified 400 conflict events in on-line journalistic news between humans and caimans within Brazilian cities between 2016 and 2021. *Caiman latirostris* (57.3%) was the most common species found in large urban centers in the Northeastern and Southeastern regions. Most encounters in which the date was informed were registered in the wet season (64.5%), and the rescues were mainly carried out by the environmental military police and fire brigades. We outlined management protocols to be undertaken by environmental agencies during crocodilian rescues to minimize animal stress and risk of accidents. The increasingly common interaction between humans and caimans in urban environments is an important alert to envisage best conducts for the coexistence between people and caimans in Brazil and shows the need to invest in public policies to mitigate the negative effects of urbanization on wildlife.

Keywords: Caimans; Habitat loss; Journalistic news; Fauna rescue, Urbanization.

1 Programa de Pós-Graduação em Biologia Animal, Universidade Federal de Pernambuco, Av. Prof. Moraes Rego, 1235 - Cidade Universitária, Recife - PE, 50670-901, Brazil.

2 Universidade Estadual Paulista, Faculdade de Ciências, Departamento de Ciências Biológicas, Bauru - SP, 17033-360, Brazil.

3 Universidade Federal do Amazonas, Departamento de Genética, Manaus - AM, 69077-000, Brazil. 4 Instituto Jacaré Conservação e Manejo de Fauna Silvestre Ltda., Rua Caçu 150, D, C62, Rio de Janeiro - RJ, 22710-074, Brazil.

5 EcoCaiman Conservação e Manejo da Biodiversidade ONG, Rua Caçu 150, D, C62, Rio de Janeiro - RJ, 22710-074, Brazil.

6 Laboratório de Caracterização de Impactos Ambientais (LCIA), Universidade Federal do Tocantins, Quadra 109 Norte, Av. NS 15, ALCNO-14, Plano Diretor Norte, Palmas - TO, 77001-090, Brazil.

7 Laboratório Vida Selvagem, Embrapa Pantanal, CP109, Corumbá, MS 79320-900, Brazil.

8 Laboratório de Ecologia Isotópica (LEI), Centro de Energia Nuclear na Agricultura/USP, Av. Centenário, 303 - Piracicaba - SP, 13400-970, Brazil.

9 Crocodylia Brasil - Grupo Brasileiro de Referência em Crocodilianos - Autoridades por Espécies e Biomas (crocodyliabrasil@gmail.com).

* Corresponding author ✉. E-mail address: PBMJ (paulobragam16@gmail.com), FM (maffei.fabio@gmail.com),

FM (fabio.lmuniz@gmail.com), RFF (institutojacare@gmail.com), TCGP (thiagoportelinha@uft.edu.br), ZC (zilca.campos@embrapa.br), LABB (luisbassetti@gmail.com)

Part of Special Issue:

Use, Management and Conservation of Wildlife in Latin America.

Edited by Hani R. El Bizri, Melina S. Simoncini, Jair H. Castro Romero, Alejandro Meléndez Herrada, Joaquín L. Navarro.

SIGNIFICANCE STATEMENT

The main objective was to present an overview of crocodylian-human conflicts in Brazilian urban areas. The urbanization process poses a real threat to crocodylian conservation due to the destruction of its natural environment. For this reason, we created standardized management protocols to help public agents responsible for the rescues in Brazilian cities.

INTRODUCTION

Cities are expanding worldwide, and urban processes have been altering landscapes and affecting ecological, biological, and behavioral processes of biodiversity in recent decades (Angel *et al.* 2012; Luniak 2004). Direct impacts (e.g., habitat fragmentation) or indirect ones (e.g., contamination of water bodies) can lead to a loss of biological diversity on a local, regional, and global scale (McDonald *et al.* 2019), including animals of various species (Vale and Prezoto 2019). Human-wildlife coexistence is a crucial goal of conservation, and the motivation for most work on conflicts has been to protect species from anthropogenic threats (Pooley *et al.* 2021a). Crocodylians consist one of the most impacted groups of vertebrates by habitat overlap with humans. In Latin American and Caribbean countries, negative interactions between humans and crocodylians have increased by 39% between 2018 and 2020 (Pooley *et al.* 2021b). This may cause fatal incidents involving humans (as documented in Colombia by Balaguera-Reina and Farfán-Ardia 2018 and in Mexico by García-Grajales and Buenrostro-Silva 2019), and destruction of natural sites due to greater anthropogenic activity (González-Desales *et al.* 2021). In Brazil, several anthropogenic activities threaten crocodylian populations, such as hydroelectric dams (Campos 2015), roads (Campos *et al.* 2012), illegal hunting (Campos and Muniz 2019; Da Silveira and Thorbjarnarson 1999; Mendonça *et al.* 2016; Mourão *et al.* 1996; Muniz *et al.* 2021) and urbanization (Mascarenhas-Júnior *et al.* 2018). Such threats mainly affect crocodylian species locally, while the natural stocks of these species at a larger scale appear not to have been substantially impacted (Campos *et al.* 2019; Magnusson *et al.* 2019; Marioni *et al.* 2021; Siroski *et al.* 2020).

Brazilian territory has the richest diversity of crocodylian species globally, with six recognized caiman species, including several distinct new evolutionary lineages (Borges *et al.* 2018; Bittencourt *et al.* 2019; Muniz *et al.* 2018; Roberto *et al.* 2020). Exponential human growth, associated with a disorderly expansion of urban areas, has resulted in huge

habitat loss, increasing the number of human conflicts (Mascarenhas-Júnior *et al.* 2018). In this modern scenario, new records are published regularly in the news and social media, reporting crocodylian occupation within cities and conflicts with local people, with a negative impact for both animals and humans, including pets or livestock (Basset 2016). Despite the increased number of cases reported in the last decade, this problematic situation has never been explored using rigorous scientific research (Balaguera-Reina and Farfán-Ardia 2018).

This manuscript aims to document crocodylian occurrence in urban areas in Brazil, creating a major database based on online local news to record and map crocodylian-human interactions in all regions across the country. Due to the complete lack of knowledge of this situation in Brazil, this database will help understand the factors responsible for increasing crocodylian-human encounters. Here we evaluated information about 1) the spatial distribution of encounters between people and crocodylian species in Brazilian urban areas, 2) the seasonality of the encounters/conflicts in each region, and 3) the possible factors that explain the increase of these encounters in some Brazilian cities. Furthermore, we outlined management protocols to be undertaken by environmental agencies during crocodylian rescues to minimize animal stress and risk of accidents, supporting government environmental agencies to create better solutions that will mitigate conflicts and help improve the management and conservation of this important group of reptiles.

MATERIAL AND METHODS

To access information on human-crocodylian interactions, we extensively searched for news reporting occurrence of crocodylians in urban areas at Google database, using the keywords “jacaré,” “resgate,” “cidade,” (that means “caiman”, “rescue”, and “city” in Portuguese), and the name of each Brazilian state, individually in quotes. We refined the searches over the past five years (01 January 2016 to 31 May 2021) using Google’s Search in “news” tab, sorting from the

most recent to the oldest, and taking care to exclude duplicate reports of the same record. Also, we activated Google alerts between January 2020 and July 2021, adding reports that were sent to us via email, with the same criteria of keywords. To minimize bias caused by Google's algorithm, all authors carried out searches between June/2021 and July/2021 in different regions of Brazil.

For each news item, when available, we filtered information related to the date, caiman species, state, city, journal name, accident events, entity responsible for the rescue/record, and the caiman release point or destination. Whenever possible, we also identified the seasonal period (dry or wet) based on the record date (Peel *et al.* 2007). In addition, to evaluate the relationship between records and season, we excluded 2021 data, which we sampled only until May, and to avoid differences between rainfall cycle across Brazilian regions. We used a chi-square test to evaluate the difference between seasons (dry and wet) and the records for each month. We considered $p < 0.05$ as our level of significance.

We used journalistic news with photos or videos to confirm the species, considering *Caiman latirostris*, *Caiman crocodilus*, *Caiman yacare*, *Paleosuchus palpebrosus*, *Paleosuchus trigonatus* and *Melanosuchus niger* as naturally distributed in Brazilian territory. Species identifications in pictures were carried out by at least two authors to ensure fidelity in identification. Only records in urban areas were considered in our sample and news with two different species were considered as one journalistic news record, but two species records. Species in reports that did not have images were considered Not Assessed (N/A). We could not determine the genus in "N/A" cases because at least two genera are distributed in the regions/states where the news without photos were published.

RESULTS AND DISCUSSION

We registered all six Brazilian crocodylian species being documented as in direct contact with humans (Figure 1A), in areas with a lot of solid waste (Figure 1B), rescued in risk areas (Figure 1C), crossing roads (Figure 1D), killed by citizens (Figure 1E), and in small lakes with dense groups (Figure 1F).

We recorded 400 journalistic news, 365 on Google Search and 35 by Google Alerts, for the last five years (2016–2021) (Figure 2). *Caiman latirostris* was the most representative species, in 57.3% (N = 229) of the total records, followed by *C. crocodilus* (18%, N = 72), *P. palpebrosus* (9%, N = 36), *M. niger* (5.5%, N = 22), *C. yacare* (5%, N = 20), *P. trigonatus* (3.8%, N = 15), and Not Assessed (1.4%, N = 6). Disregarding the Google Alerts data, most reports

(37.5%, N = 137) were published in 2020 (Figure 3), suggesting a real rising trend. Of 293 records in which date was informed, we identified that 35.5% (N = 104) occurred in the dry season, while the majority occurred in the wet season (64.5%, N = 189, $p < 0.05$) (Table 1). Aspects of crocodylian behavior and life history may account for the variation in the rate of encounters and the number of records between species in urban environments. As an example, *C. latirostris*, which has one of the largest latitudinal distributions among crocodylians (Coutinho *et al.* 2013; Verdade and Piña 2007) and easily adapts to urban environments (Freitas-Filho 2008; Marques *et al.* 2020), had the highest number of records in our study; and *P. trigonatus*, whose behavior is cryptic (Magnusson and Lima 1991), was the least recorded. Furthermore, crocodylians' behavioral patterns are often characterized by seasonality. Dry periods can reduce the movement of caimans due to reduced availability of territory and food (Campos *et al.* 2006). With the onset of the wet season, they start to disperse, searching for resources. In urban environments, rescues are mostly observed in flood periods, when water bodies are connected to floodplain areas (Negromonte *et al.* 2021), or channels (natural or artificial) connected to the urban drainage system (Oliveira *et al.* 2021), places often overlapped by roads, houses, and other buildings.

In the past two decades, the massive access to smartphones has made it possible to record wildlife, especially in urban centers (Teacher *et al.* 2013; Olson *et al.* 2014). Furthermore, the publication of records on social media has aroused the interest of the national media and websites that end up replicating these findings. Our data showed a yearly increase in news, a trend that has remained steady until the end of the study (May 2021), when in just five months, 136 events were recorded, more than 92% of the total amount reported for 2020 (N = 147). In addition to greater access to technology (e.g., smartphones and mobile Internet networks), the shutdown period during the pandemic COVID-19 contributed to reducing the use and access to streets and urban protected areas. With less people movements, the animals possibly felt safer to disperse in cities, resulting in a higher rate of encounters and records in the media (Manenti *et al.* 2020; Zellmer *et al.* 2020).

The Northeastern region of Brazil had the highest number of reports (29%, N = 116), with increased representation in the states of Pernambuco (PE; N = 36) and Bahia (BA; N = 22). The Southeast was the next (26.8%, N = 107), led by Rio de Janeiro (RJ; N = 45) and São Paulo (SP; N = 39) states. We recorded 24.8% (N = 99) of the reports for the Northern region, most in the states of Amazonas (AM; N = 33), Tocantins (TO; N = 22), and Pará (PA; N = 20).



Figure 1. *Caiman latirostris* species in urban areas. A: direct contact with humans; B: in areas with a lot of solid waste; C: rescued by researchers at neighborhoods; D: crossing roads; E: killed by citizens; F: in small lakes with dense groups.

Table 1. Percentage of journalistic news recording caimans in urban areas in different regions of Brazil during wet and dry seasons between January 2016 and May 2021. MW: Midwest; N: North; NE: Northeast; S: South; SE: Southeast.

Species	MW		N		NE		S		SE	
	Wet	Dry	Wet	Dry	Wet	Dry	Wet	Dry	Wet	Dry
<i>Caiman crocodilus</i>	85.7%	14.3%	70%	30%	47.4%	52.6%	-	-	-	-
<i>Caiman latirostris</i>	38.5%	61.5%	-	-	57.8%	42.2%	100%	0	65.8%	34.2%
<i>Caiman yacare</i>	70%	30%	-	-	-	-	-	-	-	-
<i>Melanosuchus niger</i>	-	-	65%	35%	-	-	-	-	-	-
<i>Paleosuchus palpebrosus</i>	87.5%	12.5%	70.6%	29.4%	58.3%	41.7%	-	-	-	-
<i>Paleosuchus trigonatus</i>	100%	0%	83.3%	16.7%	-	-	-	-	-	-
Not Accessed	100%	0%	100%	0%	0%	100%	-	-	-	-

On the other hand, the Midwestern region (13.3%, N = 54) was represented mainly by the states of Mato Grosso do Sul (MS; N = 28) and Mato Grosso (MT;

N = 11). Finally, the Southern region (6.1%), was represented primarily by the states of Santa Catarina (SC; N = 12) and Rio Grande do Sul (RS; N = 7)




Figure 2. Spatial distribution for human-caiman encounters recorded on digital media between 2016 and 2021. *Caiman latirostris* (star, ★), *Caiman crocodilus* (circle, ○), *Caiman yacare* (diamond, ◇), *Paleosuchus trigonatus* (triangle, △), *Paleosuchus palpebrosus* (cross, ×), and *Melanosuchus niger* (square, ◻).

(Figure 4; Additional File 1).

In the Northeast of Brazil, there are many landscapes changes and natural environment destruction in large cities, mainly in state capitals and neighboring cities, where caimans were most registered. More than 90% of the original Atlantic Rainforest has been lost to deforestation and urban expansion (Lôbo *et al.* 2011), increasing the human-crocodylian conflict by reducing caimans' habitats. The high rate of *C. latirostris* records reflects this overlap, given it is a well-distributed species in this forest domain and in urban areas in northeastern Brazil (Mascarenhas-Júnior *et al.* 2018; Siroski *et al.* 2020). This degradation scenario also occurs in other large

cities or metropolitan regions, such as Fortaleza (CE), Teresina (PI), and São Luís (MA; Silva *et al.* 2019), with a higher frequency of *C. crocodilus*.

The Northeastern region has a marked hydrological cycle with limited rainfall (up to five months) and perennial rivers (Rebouças 1997). In the Recife (PE) metropolitan region, the branched water network offers habitats and conditions for the establishment of caimans' populations (Mascarenhas-Júnior *et al.* 2018). In addition, destruction of the riparian forest exposes water bodies to roads or houses, which facilitates specimens' dispersion, and increases solid waste into the water systems (Bezerra *et al.* 2018). In Salvador (BA), the fourth largest city in Brazil,

and the main economic center in the Northeast, urbanization has been illegally occupying natural areas (Souza 2001) and weakening urban natural ecosystems (Machado *et al.* 2019). Soil and river contamination (Moura *et al.* 2010), as well as local temperature, changes due to vegetation suppression (Correia-Filho *et al.* 2019), which can directly affect the biological

cycles of urban species, mainly crocodylians, considering they are ectothermic and potential top predators in aquatic environments (Somaweera *et al.* 2020).

In Rio de Janeiro state, the recent expansion of large cities has caused the destruction of local landscapes and natural environments (such as restingas [sandbars or shoals], mangroves, and the Atlantic




Figure 3. Records for humans and caiman species published on digital media and tracked using Google searches between 2016 and 2021 in Brazil.




Figure 4. Human-caimans encounters recorded in online journalistic news in Brazil, between 2016 and 2021. Brazilian state names are abbreviated by two capital letters. AC: Acre, AL: Alagoas, AM: Amazonas, AP: Amapá, BA: Bahia, CE: Ceará, DF: Distrito Federal, ES: Espírito Santo, GO: Goiás, MA: Maranhão, MG: Minas Gerais, MT: Mato Grosso, MS: Mato Grosso do Sul, PA: Pará, PB: Paraíba, PE: Pernambuco, PI: Piauí, PR: Paraná, RJ: Rio de Janeiro, RN: Rio Grande do Norte, RO: Rondônia, RR: Roraima, RS: Rio Grande do Sul, SE: Sergipe, SC: Santa Catarina, SP: São Paulo, TO: Tocantins.

Rainforest; Díez and Montezuma 2021), which led to an increase in the occurrence of animals in urban areas. One of the most altered landscapes in Rio de Janeiro is the Lagoon Complex of Jacarepaguá, a well-known area for wild *C. latirostris* populations (Freitas-Filho 2008). Since 2010, mainly because of the choice of Rio de Janeiro as the host city of the Soccer World Cup and 2016 Olympic Games, many natural environments have been occupied (Lopes *et al.* 2014; Peres *et al.* 2018) and wetlands backfilled, drastically changing the flooded areas used by caimans. During dispersion in the search for new territories, several caimans are found in sewer channels or rescued on highways and houses (Freitas-Filho *et al.* 2013).

It is estimated that more than 70% of the Brazilian human population lives in regions covered by the Atlantic Rainforest, with more than 145 million people inhabiting 3,429 municipalities, which represents 61% of Brazilian cities (SOSMA 2021). Fragmentation of the biome for urban expansion limits access to essential resources for crocodylians, such as food, territory, and nesting areas (López-Luna *et al.* 2011). With the lack or scarcity of these resources, animals tend to disperse to new habitats, often seeking food offered by humans, creating a behavior of proximity and conditioning the caimans to settle in these areas waiting for food (Freitas-Filho *et al.* 2009). In addition, cities with many rivers, which create a branched water network, sewage systems and rainwater channels, facilitates the movement of caimans in urban environments, favoring their appearance in risk areas (Mascarenhas-Júnior *et al.* 2018).

Most of encounters in the Midwest region occurred between *C. latirostris* and people (N=12) who frequented a downtown natural lake in Três Lagoas city in the state of Mato Grosso do Sul. Individuals of *C. latirostris* can easily be spotted basking or walking on the running track that surrounds the lagoon, resulting in several encounters with people. In two events, four *C. latirostris* males (snout-vent length - SVL > 90 cm) were removed from Lagoa Maior lake and released carrying radio transmitters in the Reserva Particular do Patrimônio Natural-RPPN Cisalpina (Campos, pers.com.). The entire rescue and translocation operation was carried out by the Municipal Environmental Secretariat, under the guidance of specialists, led by Zilca Campos, who also provided advice on measures to ensure safe interaction between animals and the people who use the area around the Lagoa Maior.

One of our records indicates that species introduced outside their natural occurrence area still maintain their populations in urban centers, such as *C. yacare*, reported in the municipality of Paulínia (SP), about 500 km away from its distribution (Campos *et al.* 2020). Initially, these animals were in-

troduced into the state of São Paulo in the 1980s, with authorization from the national environmental agency, IBAMA (Instituto Brasileiro do Meio Ambiente e dos Recursos Renováveis), which allowed the installation of commercial breeding sites outside the occurrence area of the species. Most of these projects were abandoned, resulting in an incalculable number of releases in natural water bodies (Verdade 2004). The introduction of exotic or allochthonous species can be potentially harmful to ecosystems, and the presence of crocodylians outside their natural range is harmful to native species (Verdade 2004; Da Silveira *et al.* 2020). Introductions of *C. crocodilus* with established populations in South Florida, USA (Ellis 1980), and the escape of *Crocodylus moreletii* to the west coast of Mexico (Rodriguez *et al.* 2008), have been documented, potentially causing irreparable damage to local species and habitats.

The Northern and the Midwestern regions are home to the most remarkable crocodylian diversity in the country (five species), but this is not necessarily related to the higher number of human-crocodylian encounters in Brazil. Despite a high rate of urbanization, it still contains many legal protected areas around urban centers (Silva *et al.* 2017). According to our data, a large Black Caiman (*M. niger*) caused one fatal accident in a city in the state of Tocantins. Although uncommon, accidents involving this species have already been reported for the Brazilian Amazon (Bergamasco *et al.* 2018; Haddad-Júnior and Fonseca 2011; Marioni *et al.* 2013). It is noteworthy that this is the largest caiman species in the Brazilian territory (Thorbjarnarson 2010), and some of its attacks can be deadly. In the case mentioned above, the urban population killed the Black Caiman in retaliation for the incident, a practice reported for some countries in Latin America and the Caribbean (Pooley *et al.* 2021b). Over the past 20 years, Brazil has led the ranking of cases involving fatal crocodylian attacks in South America, with *M. niger* responsible for most of them (Pooley *et al.* 2021b). Those negative interactions can lead to aggressive measures from the population, such as nest destruction and crocodylian killings (Pooley 2020; Pooley *et al.* 2021a), even in situations of low risk and involving smaller species, such as species within the *Caiman* and *Paleosuchus* genera. The concern about *M. niger* occurrences in urban areas needs to be highlighted, as individuals can reach a total length of 6 meters (Medem 1983), thus representing a greater risk of fatal accidents. In addition, it is essential to emphasize that state capitals had a higher frequency of records within urban areas. Manaus is the city with the most reports (N = 28, 7% from Black Caiman), three times more cases than Belém (N = 9, 33% from Black Caiman), the second city in the North with the highest number of occur-

rences. The high number of records from Manaus may be explained by the marked urbanization process, associated with uncontrolled growth towards the preserved environments in the outskirts, the abundance of water bodies, and the forest fragments inside the city.

Crocodylian rescues were reported in 84% of journalistic news (N = 336). Most procedures were carried out by public agencies, such as fire brigades (N = 124), police (military or civilian; N = 119), and municipal environmental brigades (N = 38). On the other hand, 62 rescues (involving capture and handling) were carried out by the population and 57 by other fauna agencies that eventually helps in these procedures (e.g., Municipal Environmental Secretariats, Zoos, and Civil Defense). In Brazil, wildlife management must be carried out by authorized persons or organizations (private or public), even in urban areas. Until 2014, wildlife rescue demands in urban centers were conducted mainly by the IBAMA, environmental military police, and fire brigades (Mascarenhas-Júnior *et al.* 2018; Teixeira *et al.* 2016). Recently, as of the enactment of Complementary Law 140/2011 (Brazil 2011), fauna management has been decentralized, with the competence devolving upon each Brazilian state and/or city. Those idiosyncratic conditions lead to a lack of standardization in rescues, making it difficult for the population to know the most appropriate communication channel for help. Therefore, we recommend that environmental organizations or fauna rescue agencies should be the first to be contacted during risk situations with caimans, considering that they are trained and have adequate equipment for handling the animals. A good rescue by trained people or organizations can prevent accidents, or even animal death. Moreover, researchers and groups that work directly with caimans can provide technical support when necessary, such as the Instituto Jacaré in Rio de Janeiro-RJ (Freitas-Filho *et al.* 2013), Projeto Caiman in Vitória-ES (Mello *et al.* 2019) and Projeto Jacaré in Recife-PE (Mascarenhas-Junior *et al.* 2018).

The destination of the caiman was frequently reported (72.5%, N = 290). Returning specimens to their natural environment was the primary choice (72.4%, N = 210), followed by rehabilitation and screening centers (16.2%, N = 47), even when the individual had apparently good health conditions. In 17 cases (5.8%), caimans were relocated to other institutions (e.g., universities and zoos), but without information regarding their destination, and in 16 records (5.6%), caimans were found dead (seven of them killed by the local population). The destination of the animals after rescue is one of the most critical steps in the entire process involving conflicts with urban crocodylians. Goulart *et al.* (2010) recommend that rescued

animals should be sent to screening and rehabilitation centers for possible veterinary treatment and subsequent release (Teixeira *et al.* 2016). It is vital to collect specimen data to monitor populations during the process and determine critical points of conflict. In cases where release is possible, we recommend that caimans should be moved far from the urban center, but prioritizing the same watershed, aiming to mitigate interactions with humans and potential damage to the environment (Da Silveira *et al.* 2020).

One of the significant challenges of managing Brazilian fauna is standardizing protocols of rescue and destination specimens. The decentralization of environmental activities makes this demand difficult, but solutions must be sought. More and more professionals who work with crocodylians in Brazil are strengthening relationships with, sharing experiences with, and joining in representative groups from all regions, such as Crocodylia Brasil, a Brazilian Reference Group on Crocodylians - Authorities by Species and Biomes (Basseti *et al.* 2021). We strongly recommend that institutions responsible for fauna management (at state and municipal levels) contact crocodylian specialists so that the best procedures are carried out regarding the management of caimans in urban environments. In this sense, we developed a proposal for a standard protocol for the proper rescue, health treatment, data collection and destination of crocodylians in urban environments (Figure 5). It is important for humans to understand that wildlife is part of daily life in cities, and rescues should only be performed if animals and/or people are at risk. Under these conditions, we suggest that people contact the fire departments, military police and guards or local environmental agencies for the proper rescue. The capture and containment procedure must be carried out by a trained team with adequate equipment, such as: locking cable snares, for capture; cloths, to cover the eyes and minimize the caiman's stress; and adhesive tapes and ropes, to close and immobilize jaws and limbs, respectively, preventing bites and abrupt movements. After the process, the caiman must be taken to rehabilitation and/or screening centers for later release in a natural water body far from the city, but in the same hydrographic basin. We also suggest taking basic biometric data (at least total and snout-vent length) and biological samples, as well as conducting scutes cut, associated with tags or microchips (when possible) to mark rescued specimens, so that they can be properly monitored. This standard protocol, allied with other local recommendations from specialists in crocodylians, is important to promote animal welfare, reduce the risk of accidents, and encourage caiman protection and conservation policies within cities.




Figure 5. Flowchart of actions to be taken in rescue and translocation of crocodylians in urban areas of Brazilian cities – Training of those involved in operations to be carried out by the Municipal Environmental Secretariats.

CONCLUSION

1. *Caiman latirostris* was the most recorded species and had the widest distribution in Brazilian urban centers;

2. Encounters between humans and caimans have gradually increased in the last two years, which could be associated with access to technology, mobile cellular networks, and probably influenced by lockdown caused by the SARS-CoV-2 pandemic scenario;

3. Urban expansion and the environmental, behavioral, and natural history of crocodylians are possibly directly related to caiman occurrence and records in urban environments, especially in the wet season and in the Southeastern, Northeastern and parts of Northern regions of Brazil;

4. The contact between humans and caimans in urban areas can lead to conflicts, such as accidents (fatal or not) and pet deaths, leading to extreme measures by the population. Therefore, it is suggested that government agencies should consider crocodylian-human interaction, protection strategies, and control measures in their public policies;

5. Solid strategies and standardized management protocols for crocodylians must be implemented at the national level, including public agencies that could receive technical training from research groups and experts in the field.

ACKNOWLEDGEMENT

This publication is funded in part by the Gordon and Betty Moore Foundation through Grant No. GBMF9258 to the Comunidad de Manejo de Fauna Silvestre en la Amazonía y en Latinoamérica (COM-FAUNA). Members of Crocodylia Brasil helped with information about urban caimans.

DATA AVAILABILITY

The data used to support the findings of this study are available from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

CONTRIBUTION STATEMENT

Conceived the presented idea: PBMJ, FM, LABB, ZC.

Carried out the experiment: PBMJ, FM, FM, RFF, TCGP, LABB, ZC.

Carried out data analysis: PBMJ, FM, FM, ZC. Wrote the first draft of the manuscript: PBMJ, FM, FM, RFF, TCGP, LABB, ZC.

Review and final writing of the manuscript: PBMJ, FM, FM, RFF, TCGP, LABB, ZC.

Supervision: ZC, LABB.

REFERENCES

Angel S, Blei AM, Civco DL, Parent J (2012) **Atlas of Urban Expansion (Lincoln Institute of Land Policy)**. Lincoln Institute of Land Policy, Cambridge, MA, USA. pp. 397.

Basset LAB (2016) **Estado sanitário do jacaré-de-papo-amarelo (*Caiman latirostris*) em paisagens antropizadas no Estado de São Paulo**. PhD Thesis, Universidade de São Paulo, Piracicaba, BR.

Bassetti LAB, Da Silveira R, Botero-Arias R, Campos Z, Magnusson WE, Verdade LM, Portelinha TCG, Roberto IJ, Kluczkowski-Junior A, Marioni B, Mascarenhas-Junior PB, Muniz F, Maffei F, Araújo DD, Moura EC (2021) **Crocodylia Brazil Group (Crocbr): A Reference Network In The Research And Conservation Of Brazilian Crocodylians**. *Crocodile Specialist Group Newsletter* 39: 13-14.

Balaguera-Reina SA, Farfán-Ardila NIDIA (2018) **Are we ready for successful apex predator conservation in Colombia? Human-crocodylian interactions as a study case**. *Herpetological review* 49:5-12.

Bergamasco CJJ, Pereira RM, Rios Rodriguez JE, Fernandes Cabral BT (2018) **Thoracic trauma by black caiman's bite in the Amazon region**. *Journal of Cardiothoracic Trauma* 3:24-6.

Bezerra APXG, Gabriel FA, Lorena EMG, Santos IGS, Silva MC, Neto FCR (2018) **Environmental Diagnosis of Permanent Preservation Areas (APP) of the Capibaribe River in the city of Recife-PE**. *Revista Geama* 4:5-12.

Bittencourt PS, Campos Z, Muniz F, Marioni B, Souza BC, Da Silveira R, De Thoisy B, Hrbek T, Farias IP (2019) **Evidence of cryptic lineages within a small South American crocodylian: the Schneider's dwarf caiman *Paleosuchus trigonatus* (Alligatoridae: Caimaninae)**. *PeerJ* 7:e6580.

Borges VS, Santiago PC, Lima NG, Coutinho ME, Eterovick PC, Carvalho DC (2018) **Evolutionary significant units within populations of Neotropical broad-snouted caimans (*Caiman latirostris*, Daudin, 1802)**. *Journal of Herpetology* 52:282-288.

Brasil (2011) **Lei Complementar nº 140**. Presidência da República - Casa Civil. Brasília/DF. [http://www.planalto.gov.br/ccivil_03/leis/lcp/lcp140.htm] Accessed 13 July 2021.

Campos Z (2015) **Size of caimans killed by humans at a hydroelectric dam in the Madeira River, Brazilian Amazon**. *Herpetozoa* 28:101-104.

Campos Z, Coutinho M, Mourão G, Bayliss P, Magnusson WE (2006) **Long distance movements by *Caiman crocodylus yacare*: implications for management of the species in the Brazilian Pantanal**. *The Herpetological Journal* 16:123-132.

Campos Z, Llobet A, Magnusson WE, Piña C (2020) ***Caiman yacare* (in IUCN Red List of Threatened Species 2020)**. IUCN. e.T46586A3009881. [<https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T46586A3009881>] Accessed 18 June 2021.

Campos Z, Magnusson WE, Muniz F. (2019) ***Paleosuchus trigonatus* (in IUCN Red List of Threatened Species 2019)**. IUCN. e.T46588A3010035. [<http://dx.doi.org/10.2305/IUCN.UK.2019-1.RLTS.T46588A3010035.en>]

Accessed 20 June 2021.

Campos Z, Mourão G (1995) *Caiman latirostris* (Broad-snouted caiman). Nesting. *Herpetological Review* 26:203-204.

Campos Z, Muniz F, Magnusson WE (2012) **Dead *Paleosuchus* on roads in Brazil.** *Crocodyle Specialist Group Newsletter* 31:12-14.

Campos Z, Muniz FL (2019) **Multiple uses of Cuvier's dwarf caiman, *Paleosuchus palpebrosus*, in the semi-arid region of Northeastern Brazil.** *Crocodyle Specialist Group Newsletter* 38:5-9.

Correia-Filho WLF, Santiago DB, Oliveira-Júnior JF, Junior CAS (2019) **Impact of urban decadal advance on land use and land cover and surface temperature in the city of Maceió, Brazil.** *Land Use Policy* 87:104026.

Coutinho ME, Marioni B, Farias IP, Verdade LM, Bassetti LAB, Mendonça SH, Vieira TQ, Magnusson WE, Campos Z (2013) **Avaliação do risco de extinção do jacaré-de-papo-amarelo *Caiman latirostris* (Daudin, 1802) no Brasil.** *Biodiversidade Brasileira-BioBrasil* 1:13-20.

Da Silveira R, Botero-arias R, Marioni B, Kluczkowski-Junior A, Moura E, Maffei F, Araújo DD, Portelina TCG, Roberto IJ, Mascarenhas-Junior PB, Muniz F, Campos Z, Magnusson WE, Verdade LM, Bassetti LAB (2020) **Rearing Black Caiman (*Melanosuchus niger*) outside its distribution range concerns the brazilian crocodylian research community.** *Crocodyle Specialist Group Newsletter* 39:20-22.

Da Silveira R, Thorbjarnarson JB (1999) **Conservation implications of commercial hunting of black and spectacled caiman in the Mamirauá Sustainable Development Reserve, Brazil.** *Biological Conservation* 88:103-109.

Díez MC, Montezuma RDC (2021) **Edible Restinga: A Socio-Environmental Proposal Inside the Urban Landscape in the West Side of Rio de Janeiro, Rio de Janeiro-Brazil.** *Sociedade & Natureza* doi: [10.14393/SN-v33-2021-58037](https://doi.org/10.14393/SN-v33-2021-58037).

Ellis TM (1980) *Caiman crocodilus*: an established exotic in South Florida. *Copeia* 1980:152-154.

Freitas-Filho RF (2008) **Dieta e Avaliação de Contaminação Mercurial no Jacaré-de-Papo-Amarelo, *Caiman latirostris*, Daudin 1802, (Crocodylia, Alligatoridae) em Dois Parques Naturais no Município do Rio de Janeiro, Brasil.** MSc Thesis, Federal University of Juiz de Fora, Juiz de Fora, Brazil.

Freitas-Filho RF (2013) **Ecologia urbano do jacaré de papo amarelo (*Caiman latirostris*, Daudin 1802) no Complexo Lagunar de Jacarepaguá, Rio de Janeiro, RJ.** PhD Thesis, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil.

Freitas-Filho RF, Piña CI, Moulton TP (2009) **"Our Hidden Enemy" and the Irrational Fear of Crocodylians.** *Crocodyle Specialist Newsletter* 28:8-9.

García-Grajales J, Buenrostro-Silva A (2019) **Assessment of human–crocodyle conflict in Mexico: patterns, trends and hotspots areas.** *Marine and Freshwater Research* 70:708-720.

González-Desales GA, Sigler L, García-Grajales J, Charruau P, Zarco-González M M, Balbuena-Serrano A, Monroy-Vilchis O (2021) **Factors influencing the occurrence of negative interactions between people and crocodylians in Mexico.** *Oryx* doi: [10.1017/S0030605319000668](https://doi.org/10.1017/S0030605319000668).

Goulart VDLR, Teixeira CP, Young RJ (2010) **Analysis of callouts made relative to wild urban marmosets (*Callithrix penicillata*) and their implications for urban species management.** *European Journal of Wildlife Research* doi: [10.1007/s10344-009-0362-4](https://doi.org/10.1007/s10344-009-0362-4).

Haddad-Júnior V, Fonseca WC (2011) **A fatal attack on a child by a black caiman (*Melanosuchus niger*).** *Wilderness & environmental medicine* doi: [10.1016/j.wem.2010.11.010](https://doi.org/10.1016/j.wem.2010.11.010).

Lôbo D, Leão T, Melo FP, Santos AM, Tabarelli M (2011) **Forest fragmentation drives Atlantic forest of northeastern Brazil to biotic homogenization.** *Diversity and Distributions* doi: [10.1111/j.1472-4642.2010.00739.x](https://doi.org/10.1111/j.1472-4642.2010.00739.x).

Lopes LA, Oliveira MM, Soares CA (2014) **The effects of the 2014 World Cup in real estate market in Rio de Janeiro and São Paulo.** *International Proceedings of Economics Development and Research* doi: [10.7763/IPEDR](https://doi.org/10.7763/IPEDR).

López-luna MA, Hidalgo-mihart MG, Aguirre-león G (2011) **Description of Morelet's crocodile *Crocodylus moreletii* nests from an urban landscape in southeastern Mexico.** *Acta zoológica mexicana* 27:1-16.

Luniak M (2004) **Synurbization–adaptation of animal wildlife to urban development.** In *Proceedings 4th international urban wildlife symposium* (pp. 50-55). University of Arizona.

Machado RA, Oliveira AG, Lois-González RC (2019) **Urban ecological infrastructure: The importance of vegetation cover in the control**

of floods and landslides in Salvador/Bahia, Brazil. *Land Use Policy* 89:104180.

Magnusson WE, Campos Z, Muniz F (2019) *Paleosuchus palpebrosus* (in IUCN RedList of Threatened Species 2019). IUCN. e.T46587A3009946. [[10.2305/IUCN.UK.2019-1.RLTS.T46587A3009946.en](https://doi.org/10.2305/IUCN.UK.2019-1.RLTS.T46587A3009946.en)] Accessed 20 June 2021.

Magnusson WE, Lima AP (1991) The ecology of a cryptic predator, *Paleosuchus trigonatus*, in a tropical rainforest. *Journal of Herpetology* 25:41-48.

Manenti R, Mori E, Di Canio V, Mercurio S, Piccone M, Caffi M, Brambilla M, Ficetola GF, Rubolini D (2020) The good, the bad and the ugly of COVID-19 lockdown effects on wildlife conservation: Insights from the first European locked down country. *Biological conservation* 249:108728.

Marioni B, Barão-Nóbrega JAL, Botero-Arias R, Muniz F, Campos Z, Da Silveira R, Magnusson WE, Villamarín F (2021) Science and conservation of Amazonian crocodylians: a historical review. *Aquatic Conservation: Marine and Freshwater Ecosystems* 31:1056-1067.

Marioni B, Farias I, Verdade LM, Bassetti LAB, Coutinho ME, Mendonça SH, Veira TQ, Magnusson WE, Campos Z (2013) Avaliação do risco de extinção do jacaré-açu *Melanosuchus niger* (Spix, 1825) no Brasil. *Biodiversidade Brasileira-BioBrasil* 1:31-39.

Marques TS, Bassetti LAB, Lara NR, Portelinha TC, Piña CI, Verdade LM (2020) Home Range and Movement Pattern of the Broad-Snouted Caiman (*Caiman latirostris*) in a Silviculture Dominated Landscape. *South American Journal of Herpetology* 16:16-25.

Mascarenhas-Júnior PB, Santos EM, Correia JMS (2018) Diagnóstico dos resgates de jacarés na região metropolitana do Recife, Pernambuco. *Revista Ibero-Americana de Ciências Ambientais* 9:138-145.

McDonald RI, Mansur AV, Ascensão F, Crossman K, Elmquist T, Gonzalez A, Güneralp B, Haase D, Hamann M, Hillel O, Huang K, Kahnt B, Maddox D, Pacheco A, Pereira HM, Seto KC, Simkin R, Walsh B, Werner AS, Ziter C (2020) Research gaps in knowledge of the impact of urban growth on biodiversity. *Nature Sustainability* 3:16-24.

Medem F (1983) los crocodylia de Sur America. 2 ed. Colciencias, universidad Nacional de colombia, Bogota, pp. 259.

Mello B, Nóbrega Y, Santos M (2019) Projeto Caiman - Environmental Education. *Crocodyle Specialist Group Newsletter* 38:15-17.

Mendonça WCDS, Marioni B, Thorbjarnarson JB, Magnusson WE, Da Silveira R (2016) Caiman hunting in central Amazonia, Brazil. *The Journal of Wildlife Management* 80:1497-1502.

Moura MCS, Moita GC, Neto JMM (2010) Analysis and assessment of heavy metals in urban surface soils of Teresina, Piauí State, Brazil: a study based on multivariate analysis. *Comunicata Scientiae* 1:120-120.

Mourão G, Campos Z, Coutinho M, Abercrombie C. (1996) Size structure of illegally harvested and surviving caiman *Caiman crocodylus yacare* in Pantanal, Brazil. *Biological Conservation* 75:261-265.

Muniz F, Campos Z, Bittencourt PS, Hrbek T, Farias IP (2021) Report of poaching of dwarf caimans *Paleosuchus* spp. (Alligatoridae: Caimaninae) for meat consumption in northern Brazilian Amazon. *Herpetology Notes* 14:661-665.

Muniz F, Campos Z, Rangel SH, Martínez JG, Souza BC, De Thoisy B, Botero-Arias R, Hrbek T, Farias IP (2018) Delimitation of evolutionary units in Cuvier's dwarf caiman, *Paleosuchus palpebrosus* (Cuvier, 1807): insights from conservation of a broadly distributed species. *Conservation genetics* 19:599-610.

Negromonte MG, Mascarenhas-Junior, PB, Santos EM, Correia JS (2021) Resgate, apreensão e destinação da herpetofauna pelos órgãos de fauna na região metropolitana do Recife, pernambuco - contribuição para conservação. *Revista Ibero-Americana de Ciências Ambientais* 12.

Oliveira VCS, Viana PF, Gross MC, Feldberg E, Da Silveira R, Cioffi MB, Bertollo LAC, Schneider CH (2021) Looking for genetic effects of polluted anthropized environments on *Caiman crocodylus crocodylus* (Reptilia, Crocodylia): A comparative genotoxic and chromosomal analysis. *Ecotoxicology and Environmental Safety* 209:111835.

Olson DD, Bissonette JA, Cramer PC, Green AD, Davis ST, Jackson PJ, Coster DC (2014) Monitoring wildlife-vehicle collisions in the information age: how smartphones can improve data collection. *PloS one* 9: e98613.

Peel MC, Finlayson BL, McMahon TA (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrology and earth system sciences discussions* 4:439-473.

- Peres LF, Lucena AJ, Rotunno-Filho OC, França JRF (2018) **The urban heat island in Rio de Janeiro, Brazil, in the last 30 years using remote sensing data.** *International journal of applied earth observation and geoinformation* 64:104-116.
- Pooley S (2020) **Croc Digest: A bibliography of human-crocodile conflict research and reports.** 3 ed. Simon Pooley, London, England.
- Pooley S, Bhatia S, Vasava A (2021a) **Rethinking the study of human-wildlife coexistence.** *Conservation Biology* 35:784-793.
- Pooley S, Siroski PA, Fernandez L, Sideleau B, Ponce-Campos P (2021b) **Human-crocodylian interactions in Latin America and the Caribbean region.** *Conservation Science and Practice* 3:e351.
- Rebouças ADC (1997) **Água na região Nordeste: desperdício e escassez.** *Estudos avançados* 11:127-154.
- Roberto IJ, Bittencourt PS, Muniz FL, Hernández-Rangel SM, Nóbrega YC, Ávila RW, Souza BC, Alvarez G, Miranda-Chumacero G, Campos Z, Farias IP (2020) **Unexpected but unsurprising lineage diversity within the most widespread Neotropical crocodylian genus *Caiman* (Crocodylia, Alligatoridae).** *Systematics and Biodiversity* 18:377-395.
- Rodriguez D, Cedeño-Vázquez JR, Forstner MR, Densmore III LD (2008) **Hybridization between *Crocodylus acutus* and *Crocodylus moreletii* in the Yucatan Peninsula: II. Evidence from microsatellites.** *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* 309:674-686.
- Silva JMC, Prasad S, Diniz-Filho JAF (2017) **The impact of deforestation, urbanization, public investments, and agriculture on human welfare in the Brazilian Amazonia.** *Land Use Policy* 65:135-142.
- Silva AMC, Cunha MCC, Lopes DV (2019) **Qualidade da água como reflexo de atividades antrópicas em bacias hidrográficas do Nordeste, Brasil.** *Geosul* 34: 102-123.
- Siroski P, Bassetti LAB, Piña C, Larriera A (2020) ***Caiman latirostris* (in IUCN Red List of Threatened Species 2020).** IUCN e.T46585A3009813. [<https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T46585A3009813.en>] Accessed 18 June 2021.
- Somaweera R, Nifong J, Rosenblatt A, Brien ML, Combrink X, Elsej RM, Gordon G, Magnusson W, Mazzotti F, Percy A, Platt S, Shirley M, Tellez M, Van der Ploeg J, Webb G, Whitaker R, Webber B (2020) **The ecological importance of crocodylians: towards evidence-based justification for their conservation.** *Biological Reviews* 95:936-959.
- SOSMA (2021) **SOS Mata Atlântica.** [<https://www.sosma.org.br/conheca/mata-atlantica/>] Accessed in 13 July 2021.
- Souza AG (2001) **Favelas, invasões e ocupações coletivas nas grandes cidades brasileiras—(Re) Qualificando a questão para Salvador-BA.** *Cadernos Metrópole* 5: 63-89.
- Teacher AG, Griffiths DJ, Hodgson DJ, Inger R (2013) **Smartphones in ecology and evolution: A guide for the app-rehensive.** *Ecology and Evolution* 3:5268-5278.
- Teixeira CP, Passos L, Goulart VD, Hirsch A, Rodrigues M, Young RJ (2016) **Evaluating patterns of human-reptile conflicts in an urban environment.** *Wildlife research* 42:570-578.
- Thorbjarnarson JB (2010) **Black caiman *Melanosuchus niger*.** In: Manolis SC, Stevenson C (eds) *Crocodyles. Status survey and conservation action plan.* 3 ed. Crocodile Specialist Group, Darwin, pp. 29-39.
- Vale CA, Prezoto F (2019) **Fauna Urbana: quem vive aqui?** *CES Revista* 33:119-146.
- Verdade LM (2004) **A exploração da fauna silvestre: jacarés, sistema e recursos humanos.** *Biota Neotropica* 4:1-12.
- Verdade LM, Piña CI (2007) **O jacaré-de-papo-amarelo (*Caiman latirostris* Daudin, 1802).** *Herpetologia no Brasil* 2:295-307.
- Zellmer AJ, Wood EM, Surasinghe T, Putman BJ, Pauly GB, Magle SB, Lewis JS, Kay CAM, Fidino M (2020) **What can we learn from wildlife sightings during the COVID-19 global shutdown?** *Ecosphere* 11:e03215.

Received: 05 August 2021
Accepted: 03 November 2021
Published: 15 December 2021

Additional Files

Add File 1. Number of records of journalistic news on caimans in urban areas in different regions and states of Brazil between January 2016 and May 2021.

Region	State	City	Number of records
Midwest	DF	Brasília	7
		Goiânia	1
	GO	Itaberaí	1
		Itumbiara	1
		Nova Gama	1
		Porangatu	2
		Rio Verde	1
		São Miguel do Araguaia	1
		Aquidauana	2
	MS	Bataguassu	2
		Campo Grande	4
		Corumbá	2
		Coxim	1
		Ladário	3
		Miranda	2
		Nova Alvorada do Sul	1
		Paranaíba	1
		Porto Murtinho	1
		Três Lagoas	9
	MT	Alta Floresta	3
Cana Brava do Norte		1	
Cuiabá		6	
North	AC	Sinop	1
		Cruzeiro do Sul	2
	AM	Rio Branco	4
		Sena Madureira	3
		Careiro da Várzea	1
	AP	Iranduba	1
		Itacoatiara	2
AP	Manaus	28	
	Tapauá	1	
AP	Macapá	5	

		Belém	9
		Breves	1
		Itaituba	1
	PA	Marabá	3
		Marituba	1
		Novo Progresso	1
		Santarém	4
	RO	Monte Negro	1
		Porto Velho	3
	RR	Boa Vista	5
		Rorainópolis	1
		Araguacema	1
		Araguaína	1
		Araguatins	1
	North	Cariri	1
		Colinas	1
		Formoso do Araguaia	2
		Guaraí	1
	TO	Gurupi	3
		Lagoa da Confusão	2
		Marianópolis	1
		Miracema	1
		Miranorte	1
		Palmas	3
		Peixe	1
		Porto Nacional	1
		Silvanópolis	1
		Arapiraca	1
		Branquinha	1
		Coruripe	2
		Maceió	3
	AL	Marechal Deodoro	7
		Palmeira dos Índios	1
		Pilar	1
	Northeast	Porto de Pedras	1
		Santana do Mundaú	1
		Canudos	1
		Eunápolis	1
		Feira de Santana	1
	BA		

		Itaberaba	1
		Itamaraju	1
		Juazeiro	1
		Porto Seguro	1
		Salvador	14
		Valença	1
		<hr/>	
		Caucaia	2
		Fortaleza	1
	CE	Granja	1
		Horizonte	1
		Ipueiras	1
		Paracuru	1
		<hr/>	
	MA	São Luís	2
		Cabedelo	1
		Campina Grande	1
	PB	João Pessoa	8
		Lagoa Seca	1
		Picos	1
		Pitimbu	1
		São Bento	1
		<hr/>	
		Cabo de Santo Agostinho	2
		Camaraçibe	1
		Gameleira	1
		Igarassu	1
	PE	Jaboatão dos Guararapes	3
		Olinda	4
		Paulista	8
		Petrolina	2
		Recife	14
		<hr/>	
		Esperantina	1
		Parnaíba	1
	PI	Picos	4
		Piripiri	1
		Teresina	4
		<hr/>	
		Ceará-Mirim	1
	RN	Natal	2
		São Gonçalo do Amarante	1
		<hr/>	
	SE	Aracaju	2
		<hr/>	
		Matinhos	1
		Paranaguá	1
		<hr/>	
	PR		
South			

		Paranavaí	1
		Pontal do Paraná	1
		Umuarama	1
		Arroio do Sal	1
		Gravataí	1
	RS	Guaíba	2
		Osório	1
		Rio Grande	1
		Santa Maria	1
		Capivari de Baixo	1
		Concordia	1
		Florianópolis	4
		Garuva	1
South	SC	Indaial	1
		Jaguaruna	1
		Joinville	1
		Laguna	1
		Tijucas	1
		Guarapari	1
		Laranja da Terra	1
	ES	Serra	4
		Vitória	2
		Araçuaí	1
		Belo Horizonte	2
		Bom Despacho	1
		Carmo do Rio Claro	1
		Governador Valadares	1
		Janaúba	1
		Lagoa Santa	2
		Luz	1
		Montes Claros	2
		Pirapora	1
		Uberaba	2
		Barra do Piraí	1
		Campos dos Goytacazes	3
		Iguaba Grande	1
		Macaé	1
		Maricá	1
		Niterói	7
		Paraíba do Sul	1
	RJ		

	Paraty	1	
	Porto Real	1	
	Rio de Janeiro	21	
	São Gonçalo	2	
	São José do Imbassaí	1	
	Vassouras	3	
	Volta Redonda	1	
	<hr/>		
	Águas de São Pedro	1	
	Araras	1	
	Bertioga	1	
	Brejo Alegre	1	
	Caçapava	1	
	Cananea	1	
	Cananéia	4	
	Charqueada	2	
	Guaratinguetá	1	
Southeast	Holambra	1	
	Hortolândia	1	
	Itapetininga	1	
	Itapira	1	
	Jardinópolis	1	
	Limeira	1	
	SP	Marília	1
		Martinópolis	1
		Nova Lusitânia	1
		Osasco	1
Paulínia		2	
Paulistânia		1	
Pereira Barreto		1	
Piracicaba		1	
Presidente Venceslau		1	
Regente Feijó		1	
Ribeirão Preto	1		
Santos	5		
São Carlos	1		
Sorocaba	1		
Tupã	1		
	<hr/>		